

Dark matter signals at the LHC: forecasts from ton-scale direct detection experiments

Giorgio Arcadi,^a Riccardo Catena^a and Piero Ullio^b

^a*Institut für Theoretische Physik, Friedrich-Hund-Platz 1, 37077 Göttingen, Germany*

^b*SISSA and INFN, Sezione di Trieste, Via Bonomea, 265, 34136 Trieste, Italy*

E-mail: giorgio.arcadi@theorie.physik.uni-goettingen.de,
riccardo.catena@theorie.physik.uni-goettingen.de, ullio@sissa.it

ABSTRACT: The complementarity between dark matter searches at colliders and in underground laboratories is an extraordinarily powerful tool in the quest for dark matter. In the vast majority of the analyses conducted so far these dark matter detection strategies have been profitably combined either to perform global fits in the context of certain particle physics models (e.g. the CMSSM) or to estimate the prospects for a direct dark matter detection given the LHC potential of discovering new physics beyond the Standard Model. In this paper we propose an alternative strategy to combine direct and collider dark matter searches: employing the potential of the upcoming generation of 1-ton direct detection experiments, we show that for certain supersymmetric configurations it is possible to translate the information encoded in an hypothetically discovered direct detection signal into classes of expected signals at the LHC. As an illustrative application of our method, we show that for a 60 GeV neutralino thermally produced via resonant annihilations and identified by a 1-ton direct detection experiment, our approach allows to forecast a clearly identifiable prediction for a LHC final state involving three leptons and missing energy. The strategy presented in this paper to systematically translate a direct detection signal into a prediction for the LHC has the potential to significantly strengthen the complementarity between these two dark matter detection strategies.

Contents

1	Introduction	1
2	Determination of the benchmark points for Direct Detection	4
2.1	pMSSM setup and general constraints	4
2.2	DM relic density and spin-independent cross section	7
2.3	Benchmark points for Direct Detection	9
3	Direct detection	10
3.1	Galactic model	12
3.2	XENON1T	13
3.3	Bayesian forecasts	14
3.4	Statistical limitations	17
3.5	Mass and cross-section reconstruction	19
4	Collider analysis	21
4.1	Analysis of xen1	22
4.2	Analysis of xen2	25
4.3	Analysis of xen3	29
4.4	Analysis of xen4	34
5	Conclusions	34

1 Introduction

The nature of the dark matter (DM) component of the Universe remains unknown. One of the most popular solutions to this puzzle is the scenario in which DM is made of a beyond-Standard-Model (SM) particle which is stable, in thermal equilibrium in the early Universe and non-relativistic at chemical decoupling; the relic density for such a state scales approximately with the inverse of its pair annihilation rate into lighter SM particles and matches the abundance of DM as determined from cosmological measurements for a weak-interaction-strength coupling: this is the essence of the celebrated paradigm for weakly interacting massive particles (WIMPs) as DM candidates. On top of the fact that there are numerous extensions of the SM naturally embedding a WIMP DM state, to some extent the popularity of this framework relies on its very rich phenomenology, with a set of complementary detection strategies stemming from the framework itself on rather model-independent bases: One in general expects a tight correlation between the annihilation rate of DM in the early Universe (at a temperature of about one twentieth of the particle mass) and that in dark halos today (at relative velocities as small as 10 to 1000 km s⁻¹) giving

rise to a non-standard astrophysical source of photons, leptons or antimatter through which indirectly detect the DM particles. On the other hand, applying CP symmetry arguments to the annihilation process, one expects to produce DM states in inelastic scatterings of SM particles at high energy colliders, such as the LHC. Finally, based on crossing symmetry, one generally predicts that the weak coupling to ordinary matter in the early Universe corresponds to a small but still significant elastic scattering cross section for DM particles on ordinary matter, through which directly detect the DM WIMPs forming the halo of the Milky Way. Direct, indirect and collider searches for WIMPs have then been primary goals in the experimental effort towards the identification of the nature of DM in the latest twenty years or so, having reached today the stage when, in all three techniques, experiments have the capability to single out or exclude viable DM models.

In this paper we investigate some aspects of the complementarity between direct detection (DD) and collider searches. The goal is to investigate the potential of the upcoming generation of DD experiments, taking as reference a setup resembling the 1-ton realization of the XENON detector, and cross correlate the information encoded in a hypothetically discovered signal with the expected signals at the LHC. Modeling the coupling of DM and SM states at an effective operator level, such as in the recent analysis of, e.g., [1] one can drive a direct link between the two detection strategies; on the other hand, when considering explicit WIMP models cross symmetry arguments rarely apply, with the couplings ensuring the appropriate relic density most often not corresponding to those setting the WIMP-nucleus elastic scattering cross sections and those responsible for the production of DM states in LHC's pp collisions. We choose then here to focus on a definite scenario, assuming that the DM particle is the lightest neutralino in the minimal supersymmetric extension to the SM (MSSM). This choice is motivated by the fact that, with a number of degrees of freedom which is more than twice those of the SM and the large number of free parameters involved (we will refer to a simplified version of the so-called phenomenological MSSM, or pMSSM, allowing for 12 free parameters), this framework is general enough to embed several different mechanisms to provide the correct relic abundance and/or a scattering rate at a detectable level. At the same time, the neutralino is the most closely investigated WIMP DM candidate within the most widely studied framework for searches for physics beyond the SM at colliders (as well as in rare flavor processes): we will then exploit the several well-known patterns for this DM state as criteria to explore the complex parameter space and interpret results, as well refer to a set of well-established search channels to extract guidelines which are actually more general than for the MSSM itself.

Our analysis starts with the choice within the MSSM of a set of few benchmark models representative of WIMP DM candidates which, in the plane spin-independent elastic scattering cross section per nucleon versus DM mass, appear below the region excluded by DD searches – with the most relevant limit stemming from the null result recently reported by the XENON100 experiment [2] – but still giving a sizable number of scattering events for a 1-ton detector. All benchmarks are selected checking that the lightest Higgs boson in each model matches the Higgs-like state recently discovered at the LHC, and fulfilling current constraints on beyond-SM states from rare process like $b \rightarrow s\gamma$ and $B_s \rightarrow \mu^+\mu^-$ as well as general trends from direct SUSY searches at colliders. For each benchmark, first a

set of mock data for the recoil spectrum in a 1-ton detector is simulated; such spectrum depends, on top of the spin-independent scattering cross section and the DM mass, on the phase space distribution of DM particles. Uncertainties on the latter have been recently investigated in Refs. [3, 4], implementing a large set of dynamical observables for the Milky Way and under the simplifying (but reasonable) approximations of spherical symmetry and isotropy; we will exploit results of such analysis in our approach here (complementary methods to study the local DM distribution have been discussed for instance in Refs. [5–10]). Our simulated DD data samples consist in a number of measured recoil energies varying from approximately 30 to 190, depending from the benchmark model under analysis. Before analyzing these data, a statistical test has been performed on the simulated data samples to assess whether they provide a faithful realization of their true underlying distribution. This aspect has been addressed with great care since statistical fluctuations in the distribution of the observed recoil energies can potentially affect the DM mass and scattering cross section determinations significantly, in particular for heavy DM candidates with low scattering cross sections [11]. Mock data are then analyzed employing a Bayesian approach supported by a Markov Chain Monte Carlo (MCMC) Likelihood scanning technique of the parameter space defined by both the particle physics model parameters and the parameters related to the DM phase space distribution function. The results of this analysis regarding the preferred values for the DM mass and scattering cross section will be presented in terms of Bayesian credible regions. Having derived through this procedure the accuracy within which the relevant DM properties, namely mass and scattering cross section, are reconstructed and their mapping on the MSSM parameter space, we take this input to outline the most suitable collider search strategies and argue next future detection prospects. This second part of our analysis has been performed by focusing on the regions of the DM parameter space characterized by the highest statistical weight (i.e. with the largest Bayesian posterior probability density resulting from our DD analysis) and simulating for a subset of models, sampled within these regions, the ATLAS detector response in its configuration with 7 TeV center of mass energy. This allowed us, in particular in the case of one of the benchmark investigated here, to effectively translate the simulated DD signals studied in the first part of this work into definite predictions for certain LHC observables. This procedure defines a method to efficiently exploit the upcoming generations of 1-ton DD detectors to forecast valuable predictions for the LHC and it constitutes the main achievement of this paper. The paper is organized as follows: in section 2 we provide a description of the benchmarks studied in this paper and review the criteria which brought to their selection. Section 3 is instead devoted to a study of the capability of a ton-scale DD experiment in reconstructing mass and spin-independent cross section of the DM candidates associated with these benchmarks. In section 4 we will translate these informations on the DM mass and scattering cross section into a prediction for certain classes of LHC final states while our conclusions will be summarized in section 5.

2 Determination of the benchmark points for Direct Detection

This section is devoted to the determination of the benchmark points for the DD analysis. We will start by introducing the MSSM considered in this work and then we will review the dependence of the scattering cross section from the SUSY parameters stressing the implications in our framework.

2.1 pMSSM setup and general constraints

For our investigation of the MSSM parameter space we will pursue a phenomenological approach adopting the framework dubbed pMSSM [12]. Its most general realization features 22 parameters defined at the Electro-Weak (EW) scale:

- M_1 , M_2 and M_3 parameters corresponding to the gaugino masses;
- μ , m_A and $\tan\beta$ parameters describing the Higgs/Higgsino sector;
- sfermion soft-SUSY breaking mass parameters $m_{\tilde{q}}$, $m_{\tilde{u}_R}$, $m_{\tilde{d}_R}$, $m_{\tilde{l}_L}$, $m_{\tilde{e}_R}$, equal for first and second generation ;
- trilinear soft terms A_u , A_d and A_e equal for the first and second generation;
- third generation soft mass parameters $m_{\tilde{q}_{3L}}$, $m_{\tilde{l}_R}$, $m_{\tilde{b}_R}$, $m_{\tilde{l}_{3L}}$, $m_{\tilde{\tau}_R}$;
- third generation trilinear soft terms A_t , A_b and A_τ ;

with the further assumption that no additional sources, with respect to the SM, of flavor and CP violation are present. The exploration of such large parameter space is, however, computationally very demanding; for this reason we have considered a few limits reducing its dimensionality, paying attention however to still cover all the scenarios that are, possibly, phenomenologically relevant.

First of all we have neglected contribution to all observables from squarks of the first two families fixing their mass at about a few TeV, decoupled from the rest of the spectrum. In addition, their corresponding trilinear A -terms have been set to zero. This choice is motivated by the fact that current null results of SUSY searches at LHC constraint more severely masses of the squarks of the first two generations, as well as the gluino mass, ruling out values even about 1 TeV [13, 14]¹². For the same reason we have considered only values of the gluino mass greater than 600 GeV.

For simplicity we have also further reduced the parameter space in the sfermion sector by assuming $m_{\tilde{l}} = m_{\tilde{l}_L} = m_{\tilde{e}_R}$, $m_{\tilde{l}_3} = m_{\tilde{l}_{3L}} = m_{\tilde{\tau}_R}$, and $m_{\tilde{q}_{3R}} = m_{\tilde{b}_R} = m_{\tilde{t}_R}$. We have finally also set $A_b = A_\tau$.

¹Most of the exclusion limits available in the literature rely on oversimplified models, mostly on the CMSSM, while their translation into a multi-parameter model like the pMSSM is not straightforward. In the scans considered in the next subsection we will not apply excessively tight constraint determining more accurate limits case by case through our detector simulations.

²Customarily considered collider searches can be evaded in the case of spectra enforcing particularly low mass splitting among particles involved in the production as subsequent decay chains, thus weakening exclusion limits even for first family squarks. We will neglect however this possibility in order to avoid dealing with a parameters space with an excessively high dimensionality.

Given this setup we have performed a series of flat scans in order to determine the relevant scenarios for the Monte Carlo study. The 12 remaining SUSY parameters have been varied over the following ranges:

$$\begin{aligned}
10 &\leq M_1 \leq 3000 \text{ GeV} \\
100 &\leq M_2, \mu \leq 3000 \text{ GeV} \\
600 &\leq M_3 \leq 3000 \text{ GeV} \\
100 &\leq m_{\tilde{q}_{3L}}, m_{\tilde{q}_{3R}}, m_{\tilde{l}}, m_{\tilde{l}_3} \leq 3000 \text{ GeV} \\
90 &\leq m_A \leq 3000 \text{ GeV} \\
3 &\leq \tan \beta \leq 60 \\
-9 &\leq A_t, A_b \leq 9 \text{ TeV}
\end{aligned} \tag{2.1}$$

SUSY spectra have been generated with the package SUSPECT [15] and required to be physical, i.e to satisfy the EW breaking condition, and to have a neutralino as LSP. We have in addition considered a series of general bounds.

We have first of all required that the generated spectra provide a CP-even Higgs boson mass in agreement with the recent LHC discovery [16] of a neutral boson of mass of around 125.5 GeV and couplings compatible with the ones of the SM Higgs boson (there are slight indications, not statistically decisive, of some deviations from the standard values, see for details [17, 18]). The MSSM setup most likely in agreement with this result is the so-called decoupling regime [19] in which the light CP-even Higgs state h has SM-like couplings and is required to have mass in agreement with the experimental determination, while all the other Higgs states are degenerate in mass and lie at a sensitively higher scale. In this regime the Higgs mass m_h is approximatively given by the sum of tree-level term and stop-top loop contributions:

$$m_h \approx m_Z \cos 2\beta + \frac{3G_F}{\sqrt{2}\pi^2} m_t^4 \left(\log \left(\frac{M_s^2}{m_t^2} \right) + \frac{X_t^2}{M_s^2} \left(1 - \frac{X_t^2}{12M_s^2} \right) \right) \tag{2.2}$$

where $X_t \equiv A_t - \mu \cot \beta$, namely the stop mixing parameter, and $M_s = \sqrt{m_{\tilde{t}_1} m_{\tilde{t}_2}}$ with $\tilde{t}_1$ and $\tilde{t}_2$ being the two stop mass eigenstates. Matching the experimental value of the Higgs mass is rather challenging for the MSSM. Indeed it requires a large contribution from loop-corrections which achieve the correct value only in the case the stop mass scale lies above around 3 TeV unless the stop mixing parameter $X_t = A_t - \mu \tan \beta$ fulfills the so called maximal mixing condition [20, 21], namely $X_t \approx \sqrt{6}M_s$ ³. In this work we have imposed the requirement $123 \leq m_h \leq 129$ GeV, taking into account the experimental uncertainties as well as the ones affecting the numerical computation.

We just mention that correct values of the Higgs mass can be also achieved by configurations apart from the decoupling regime, featuring light, i.e. close in mass to the LHC candidate, masses for the other Higgs states (see e.g. [23] for a review). However these are

³Milder constraints are obtained in the minimal extension of the MSSM, the so called NMSSM, in which the Higgs couples with an additional superfield which provides an additional one-loop contribution to its mass [22].

severely constrained by the searches of CP-odd Higgs decays into tau pairs [24, 25] and, as we will clarify in the following, by the limits from $Br(B_s \rightarrow \mu^+ \mu^-)$. We finally notice that light DM scenarios may influence Higgs phenomenology allowing for the decay $h \rightarrow \chi\chi$.

The invisible decay width of the Higgs has been tested by some early studies based on multijet + missing E_T events originated by the associated production of the Higgs with a gauge boson or jet. This kind of studies put an upper bound on the invisible branching ratio of the Higgs of the order of 0.5 [1]. More recent studies are aiming to infer the viability of a non zero $Br(h \rightarrow \chi\chi)$ by fitting the current data relying on Higgs searches. A study of this type can be found e.g. in Ref. [26] in which a fit of the current data has been performed by adding to the decay channels into visible states a branching ratio into two DM particles made free to vary. The outcome of such analysis is that $BR(h \rightarrow \chi\chi) > 0.4$ is excluded at the 95% confidence level with the best fit pointing towards values close to zero.

The second set of general bounds applied in our analysis refers to the searches of FCNC processes, focussing on the three most important observables, namely the anomalous magnetic moment a_μ and the decay processes $b \rightarrow s\gamma$ and $B_s \rightarrow \mu^+ \mu^-$. Regarding the anomalous moment of the muon, the SUSY contributions come from neutralino-slepton loops and chargino-sneutrino loops [27] with the latter typically dominating [28]. In absence of extra sources of flavor violation the most important parameter turns out to be the mass scale of the sleptons. Being this not directly related to DD we will not consider in the scans its experimental value but rather only an upper bound, together with the condition $\mu > 0$ which enforces the preferred sign of the SUSY contribution.

Contributions to the $b \rightarrow s\gamma$ processes are instead induced by charged Higgs-top loops and chargino-stop loops [29] which have to be combined with the SM contribution, $Br_{SM}(b \rightarrow s\gamma) \simeq 3.0 \times 10^{-4}$. While the charged Higgs-top loops always add constructively to the SM contribution, the behavior of chargino/stop loops depends on the sign of the quantity $Re(\mu A_t)$. Having fixed the μ parameter to a positive value, they add destructively to the SM one when $A_t < 0$ and constructively when $A_t > 0$. We also remark that the chargino-stop loops contribution is enhanced at high $\tan\beta$.

Finally, regarding $B_s \rightarrow \mu^+ \mu^-$, the additional contributions with respect to the SM arise from neutral heavy Higgs loops and are proportional to $\tan^3\beta$ and to $A_t\mu$. This dependence on $\tan^3\beta$ and the heavy Higgs mass makes, as already stated, this process also relevant for the Higgs phenomenology. In our analysis we made use of the numerical package SuperIso [30, 31] for computing the rates of these three process imposing the range $[2.77, 4.37] \times 10^{-4}$ for the values of $Br(b \rightarrow s\gamma)$ and upper bounds of 4.79×10^{-9} for δa_μ and 4.5×10^{-9} for $Br(B_s \rightarrow \mu^+ \mu^-)$ [32, 33].

From the flat scans of the pMSSM parameter space performed varying the 12 parameters previously introduced and accounting for all the mentioned experimental bounds, we selected four benchmark points for the Monte Carlo analysis on which our strategy to combine the DD technique and DM searches at the LHC is based. Before introducing these benchmark points, in the next section we will describe which DM configurations provide cosmologically viable scenarios in light of the recent DD limits from the XENON100 experiment (and current limits on SUSY searches) and how the DM spin-independent (SI) scattering cross section depends from the underlying parameters. The main quantities for

this study have been numerically computed through the package DarkSUSY [34].

2.2 DM relic density and spin-independent cross section

The limit on the neutralino relic density is satisfied in rather different regimes, depending from whether the lightest neutralino is higgsino- and wino-like, or bino-like. In the former case, since higgsino and winos have unsuppressed couplings to W bosons via their charged counterparts, the pair annihilation rate is very large and the relic density matches the DM density in the Universe only going to the heavy mass regime, at about 1.4 and 2.7 TeV for, respectively, a pure higgsino and a pure wino [35]. Such regime (as much as for the couplings) is however not favorable for a DD analysis and will be discarded here.

Moving to bino-like DM, we have an opposite situation, since bino-like DM couples to fermions only but the s-wave pair annihilation rate is helicity suppressed, namely proportional to the square of the mass of the final state fermion. As a consequence the value of the DM relic density exceeds the experimental limit with the exception of some peculiar regions of the parameter space. Interestingly these regions typically enforce definite relations among the SUSY parameters which can be used in correlating an hypothetical DD signal with the collider detection. Some of these configurations are already well known in the context of CMSSM/SUGRA, thus for simplicity we will refer to them in the jargon borrowed from this class of models.

Bino-like DM can be cosmologically viable in the following cases:

- in presence of at least one rather low mass sfermion, namely with mass of the order of 100-200 GeV, in such a way the annihilation cross section into fermion pairs matches the required value for a thermal WIMP. This is customarily dubbed bulk region. For the SUGRA/CMSSM it is ruled out by current LHC limits while in the pMSSM is still partially viable the possibility of light sleptons as opposed to higher mass squarks in agreement with the LHC bounds.
- in the so called Higgs funnel region, in which there is an enhancement of the annihilation process, due to a s-channel resonance. In the CMSSM this can happen with the CP-odd Higgs, when $m_A \simeq 2m_\chi$. In the pMSSM it can also occur, at light DM masses, because of resonances associated to the Z boson and to the light CP-even Higgs.
- in case the DM is a mixture of a bino and a small higgsino or wino component. The first case also occurs in the CMSSM in the so called focus point region while the pMSSM admits both the two configurations which are more generically dubbed as well-tempered neutralino [36].
- in the so called coannihilation scenarios, namely when one or more particles almost degenerate in mass with the DM contribute to keeping the DM in chemical equilibrium in the Early Universe. We also notice that coannihilating particles evade most of the current SUSY searches because the low mass splittings disfavor the detection of their decay products, after an eventual production at collider.




Figure 1: Curves at the same values of the DM bino (higgsino) fraction G_g (Z_g), as reported in the plot, in the plane (M_χ, σ_{SI}) . These curves have been obtained by considering a simplified model featuring only M_1 and μ as free parameters while all the remaining mass spectrum is fixed at about few TeV. In addition m_h and $\tan \beta$ have been fixed to, respectively, 126 GeV and 10. The solid and the dashed black line are, respectively, the XENON100 limit and the XENON1T projected sensitivity.

Moving to direct detection, looking at the general form for the neutralino-quark effective coupling in the small momentum transfer limit, reported e.g in [37]⁴, we can identify two main kinds of contributions determining the SI cross section. A first contribution depending on the SUSY parameters involved in the Higgs sector comes from the t-channel interaction mediated by a CP-even Higgs. In the decoupling regime only the light Higgs state h is relevant with the coupling scaling with the gaugino-higgsino mixing being zero for a pure bino or a pure higgsino. This effect can be seen in Fig. 1, where, in a simplified model featuring only M_1 and μ as free parameters, we show the isocontour for the bino (higgsino) fraction G_g (Z_g) in the plane DM mass versus SI cross section.

The second kind of contribution comes from interactions mediated by squarks which is thus suppressed, compared to the former, by their mass scale. There is however a possible exception occurring in case of a very low mass splitting between a squark and the DM particle. Remarkably this setup suggests an interesting correlation with coannihilation scenarios. Moreover a suitable enhancement of the squark mediated interactions requires a sizable mixing between left and right-handed states. In our framework this condition can be fulfilled by third family squarks⁵. This is one relevant example of how the interplay of different observables may enforce correlations among SUSY parameters. On the other

⁴The expression there reported may not be strictly valid in case of coannihilations and need to be eventually corrected according to [38, 39].

⁵This further motivates our choice of limiting to third generation squarks the possibility of being at low mass




Figure 2: The four benchmark points studied in this paper are shown together with the present Xenon exclusion limit and an estimation for the expected sensitivity of the Xenon experiment in its 1-ton configuration.

hand the parameters of the stop mass matrix (related in our setup also to sbottom masses) determine as well the quantities X_t and M_s entering in eq. (2.2). As already mentioned the constraint on the Higgs mass favors stops (and sbottoms as well) far from the coannihilation regime with the exception of a rather narrow range of values of $|X_t/M_s|$ [21]. At the same time also the bounds from $b \rightarrow s\gamma$ seriously impact this scenario.

2.3 Benchmark points for Direct Detection

We now introduce four benchmark points selected for being representative of different correlation patterns between DD and collider observables. The first benchmark, labeled by **xen1** has been designed for the study of light DM scenarios and features a DM mass close to 56 GeV thus lying around the light CP-even Higgs resonance. We have then defined the point **xen2**, featuring DM-slepton coannihilations, with the aim of investigating the collider signals possibly induced by light sleptons. Finally, we have considered two benchmark points, namely **xen3** and **xen4**, corresponding to scenarios where DM is thermally produced through coannihilations with a light squark. The first of them, namely **xen3**, has been designed to have a very low value of the SI cross section and a mass close to the current exclusions limits, including the bounds from direct SUSY searches at LHC. The benchmark point **xen4** is instead the heaviest benchmark considered in this paper, having a mass close to 500 GeV. The main features of the four benchmarks are summarized in Tab. 1 while Fig. 2 shows the four benchmarks in the plane DM mass versus SI cross section together with the present Xenon exclusion limit and an estimation for the expected sensitivity of the Xenon experiment in its 1-ton configuration.

benchmark	m_χ	m_A	μ	$m_{\tilde{t}_1}$	$m_{\tilde{b}_1}$	$m_{\tilde{g}}$	$m_{\tilde{\tau}}$	σ_{SI}	N
xen1	56	890	291	557	735	815	165	$2.3 \times 10^{-45} \text{cm}^2$	170
xen2	155	1550	353	1715	1777	2254	163	$3.12 \times 10^{-45} \text{cm}^2$	129
xen3	194	2900	1500	340	209	2278	732	$6.1 \times 10^{-46} \text{cm}^2$	24
xen4	477	2780	718	527	673	665	656	$1.35 \times 10^{-44} \text{cm}^2$	187

Table 1: Summary table of the benchmark points. In the last column we report the number of recoil energies simulated in the DD analysis of section 3.

Starting from these benchmark points we will illustrate how following our approach an hypothetical signal in the DD technique can be translated into a prediction for certain LHC observables.

3 Direct detection

In this section we will investigate the capability of a ton-scale direct detection experiment in reconstructing mass and SI scattering cross section associated with the pMSSM benchmarks previously introduced. Direct detection experiments seek to measure the nuclear recoil energy deposited in an underground detector when an incident χ particle of the Milky Way DM halo interacts with a nucleus of the target material. In the case of spin-independent χ -nucleus interactions the differential cross section for this process can be written as

$$\frac{d\sigma}{dQ} = \sigma_{SI} \frac{M_N}{2 M_n^2 |\mathbf{u}|^2} A^2 \mathcal{F}_N^2(Q), \quad (3.1)$$

where Q is the energy transferred to the nucleus of mass M_N from a χ particle moving with velocity $\mathbf{u}$ in the detector rest frame and A is the nucleus mass number. The form factor $\mathcal{F}_N(Q)$ accounts for the internal structure of the nucleus while σ_{SI} is the χ -nucleon SI scattering cross section at zero momentum transfer and $M_n = M_\chi m_p / (M_\chi + m_p)$ the χ -nucleon reduced mass. The differential rate of scattering events per unit detector mass in a direct detection experiment is given by the convolution of the incident flux of χ particles with the differential cross section (3.1):

$$\frac{dR}{dQ} = \frac{\sigma_{SI}}{2 M_\chi M_n^2} A^2 \mathcal{F}_N^2(Q) \int_{|\mathbf{u}| > v_{\min}} d^3u \frac{\rho_\chi \cdot f_\chi(\mathbf{u}, t)}{|\mathbf{u}|}, \quad (3.2)$$

where the lower limit in the integral is the minimum velocity required for a χ particle to deposit the energy Q . In the case of an elastic scattering, such as for WIMP-nucleus interactions, the minimum velocity is given by

$$v_{\min} = \sqrt{\frac{QM_N}{2M_r^2}}. \quad (3.3)$$

In Eq. (3.3) $M_r = M_\chi M_N / (M_\chi + M_N)$ is the χ -nucleus reduced mass while M_χ represents the mass of the χ particle.

The product $\rho_\chi \cdot f_\chi \equiv F_h$ of the local DM density ρ_χ and velocity distribution f_χ is equal to the phase-space density of the halo DM particles F_h evaluated at the detector and in its reference frame. The local χ phase-space density is computed assuming that the χ particles account for the entire DM component and applying Eddington's inversion formula [40] to the mass model for the Milky Way introduced in section 3.1. This model is characterized by a spherically symmetric and isotropic distribution of DM particles and it furthermore assumes spherical symmetry for the underlying Galactic gravitational potential $\Phi(r)$. In this limit, Eddington's formula [40] gives an one-to-one correspondence between the DM halo density profile $\rho_h(r)$ and its phase-space density F_h :

$$F_h(\mathcal{E}) = \frac{1}{\sqrt{8\pi^2}} \left[\int_0^\mathcal{E} \frac{d^2\rho_h}{d\Psi^2} \frac{d\Psi}{\sqrt{\mathcal{E}-\Psi}} + \frac{1}{\sqrt{\mathcal{E}}} \left(\frac{d\rho_h}{d\Psi} \right)_{\Psi=0} \right], \quad (3.4)$$

where we introduced the relative potential $\Psi(r) = -\Phi(r) + \Phi(r = R_{vir})$ and the virial radius R_{vir} , corresponding to the radius at which the DM halo is truncated. Under the assumptions considered here, F_h is a function of the relative energy $\mathcal{E} = -E + \Phi(r = R_{vir}) = -E_{kin} + \Psi(r)$ only, with E and E_{kin} , respectively, the total and kinetic energies. The relative energy depends on the DM particle velocity and position in the Galaxy and therefore determines the space-time dependence of F_h itself. To numerically compute F_h at the local Galactocentric distance it is actually simpler to implement Eq. (3.4) by changing the integration variable from Ψ to the radius of the spherical system r . Then, the radial dependence of ρ_h and Φ from the local position in the Galaxy out to R_{vir} is the only information needed to calculate F_h . Further details on the implementation and validity of this approach can be found in [4]. In summary, Eddington's inversion formula allows us to express the integrand of Eq. (3.2) directly as

$$\rho_\chi \cdot f_\chi(\mathbf{u}, t) = F_h[\mathcal{E}_0(\mathbf{v})], \quad (3.5)$$

with $\mathbf{v}(t) = \mathbf{v}_{obs}(t) + \mathbf{u}$, being $\mathbf{v}_{obs}$ the velocity of the detector in the Galactic frame, where F_h is computed, and $\mathcal{E}_0$ the relative energy at the local Galactocentric distance. For $\mathbf{v}_{obs}$ we take the expression

$$\mathbf{v}_{obs}(t) = \mathbf{v}_{LSR} + \mathbf{v}_{\odot,pec} + \mathbf{V}_\oplus(t), \quad (3.6)$$

where the Local Standard of Rest velocity $\mathbf{v}_{LSR}$ (namely the rotation of the Sun with velocity Θ_0 around the Galactic center; calculable given a mass model for the Galaxy), the Sun's peculiar velocity $\mathbf{v}_{\odot,pec}$ and the motion of the Earth around the Sun $\mathbf{V}_\oplus$ are given by

$$\begin{aligned} \mathbf{v}_{LSR} &= (0, \Theta_0, 0), \\ \mathbf{v}_{\odot,pec} &= (U_\odot, V_\odot, W_\odot) \simeq (10, 5.2, 7.2) \text{ km/s} \\ \mathbf{V}_\oplus(t) &= V_\oplus [\hat{\varepsilon}_1 \cos \omega(t - t_1) + \hat{\varepsilon}_2 \sin \omega(t - t_1)], \end{aligned} \quad (3.7)$$

with $V_\oplus = 29.8$ km/s and $t_1 = 0.218$ is the fraction of the year before the Spring equinox. Finally, $\hat{\varepsilon}_1$ and $\hat{\varepsilon}_2$ are the directions of the Earth's velocity at times t_1 and $t_1 + 0.25$ years. In Eq. (3.7) all velocities are expressed in Galactic coordinates, where $\hat{\mathbf{x}}$ is the direction to the Galactic Center, $\hat{\mathbf{y}}$ the direction of disk rotation and $\hat{\mathbf{z}}$ identifies the North Galactic Pole.

3.1 Galactic model

The Galactic mass model adopted in the present analysis to compute F_h has been extensively investigated in Refs. [3, 4]. For completeness, we briefly summarize it in what follows. The model consists in two luminous mass components, namely the stellar disk and the Galactic bulge, and in a DM halo. The Galactic model parameters introduced in this section will be then treated as nuisance parameters in the Bayesian analysis of our pMSSM benchmarks performed in the next subsections.

Regarding the stellar disk, we assume a mass density profile which in cylindrical coordinates (R, z) with origin in the Galactic center is given by

$$\rho_d(R, z) = \frac{\Sigma_d}{2z_d} e^{-\frac{R}{R_d}} \text{sech}^2\left(\frac{z}{z_d}\right) \quad \text{with} \quad R < R_{dm}, \quad (3.8)$$

where Σ_d is the central disk surface density, R_d and z_d are length scales in the radial and vertical directions, while R_{dm} is the truncation radius of the disk. R_{dm} is assumed to scale with the local Galactocentric distance R_0 according to the prescription $R_{dm} = 12[1 + 0.07(R_0 - 8 \text{ kpc})]$ kpc and the vertical scale z_d is fixed to the best fit value suggested in Ref. [41], $z_d = 0.340$ kpc.

The bulge/bar region is instead characterized by the mass density profile:

$$\rho_{bb}(x, y, z) = \bar{\rho}_{bb} \left[s_a^{-1.85} \exp(-s_a) + \exp\left(-\frac{s_b^2}{2}\right) \right] \quad (3.9)$$

where

$$s_a^2 = \frac{q_b^2(x^2 + y^2) + z^2}{z_b^2} \quad \text{and} \quad s_b^4 = \left[\left(\frac{x}{x_b}\right)^2 + \left(\frac{y}{y_b}\right)^2 \right]^2 + \left(\frac{z}{z_b}\right)^4. \quad (3.10)$$

We implement in this analysis an axisymmetrized version of Eq. (3.9), and assume $x_b \simeq y_b = 0.9 \text{ kpc} \cdot (8 \text{ kpc}/R_0)$, $z_b = 0.4 \text{ kpc} \cdot (8 \text{ kpc}/R_0)$ and $q_b = 0.6$. See also [3] concerning the choice of these parameters. Rather than using the two mass normalization scales Σ_d and $\bar{\rho}_{bb}$ as free parameters, we re-parameterize these in terms of two dimensionless quantities, namely, the fraction of collapsed baryons f_b and the ratio between the bulge/bar and disk masses Γ :

$$f_b \equiv \frac{\Omega_{\text{DM}} + \Omega_b}{\Omega_b} \frac{M_{bb} + M_d + M_{\text{H}_I} + M_{\text{H}_2}}{M_{\text{vir}}} \quad (3.11)$$

$$\Gamma \equiv \frac{M_{bb}}{M_d}. \quad (3.12)$$

In Eq. (3.11) we also included the sub-leading contributions to the total virial mass M_{vir} (defined in the following) associated with the atomic (H_I) and the molecular (H_2) Galactic gas layers, with profiles as given in [42]. In summary, the free parameters describing the luminous components are R_0 , R_d , f_b and Γ .

Concerning the DM halo component we consider an Einasto profile [43, 44], which is favored by the latest N-body simulations and is given by

$$\rho_h(r) = \rho' f_E(r/a_h), \quad (3.13)$$

with

$$f_E(x) = \exp \left[-\frac{2}{\alpha_E} (x^{\alpha_E} - 1) \right]. \quad (3.14)$$

The reference normalization ρ' and the scale radius a_h in Eq. (3.13) are often rewritten as a function of the virial mass M_{vir} and of the concentration parameter c_{vir} by inverting the relations:

$$M_{vir} \equiv \frac{4\pi}{3} \Delta_{vir} \bar{\rho}_0 R_{vir}^3 = \frac{\Omega_{DM} + \Omega_b}{\Omega_{DM}} 4\pi \int_0^{R_{vir}} dr r^2 \rho_h(r) \quad (3.15)$$

$$c_{vir} \equiv R_{vir}/r_{-2} \quad (3.16)$$

where the virial overdensity Δ_{vir} in the first equation is computed according to Ref. [45] while $\bar{\rho}_0$ is the mean background density today. The presence in this equation of Ω_{DM} and Ω_b , the DM and baryon energy densities in units of the critical density, reflects our assumption that only a fraction equal to $\Omega_{DM}/(\Omega_{DM} + \Omega_b)$ of the total virial mass consists of DM. Their values have been set according to the mean values from the fit of the 7-year WMAP data [46]. In the second equation, instead, r_{-2} is the radius at which the effective logarithmic slope of the DM profile is equal to -2 . Finally, we assume that the baryons which do not collapse in the disk are distributed according to the same profile of the DM component. The free parameters describing the DM halo are therefore M_{vir} , c_{vir} and α .

The Galactic model used to compute F_h also includes an additional parameter, the so-called anisotropy parameter $\beta \equiv 1 - \sigma_t^2/\sigma_r^2$, where σ_r and σ_t are respectively the radial and tangential velocity dispersions of the halo stars. It has been introduced in Ref. [3] to include in the parameter estimation the velocity dispersion measurements of Ref. [47].

3.2 XENON1T

In a real experiment Eq. (3.2) has to be modified in order to account for experimental limitations related to the efficiency of the detector, its finite energy resolution and the presence of an experimental noise. In our discussion of the capability of a ton-scale direct detection experiment in reconstructing the DM mass and SI scattering cross section we will implement the experimental limitations which would apply, as a first approximation, to the XENON1T experiment. We will therefore consider a constant efficiency $\epsilon = 0.3$ and include finite energy resolution effects by assuming a Gaussian probability

$$\xi(E, Q) = \frac{1}{\sqrt{2\pi\sigma_{Xe}^2(Q)}} e^{-(E-Q)^2/2\sigma_{Xe}^2(Q)} \quad (3.17)$$

that to an event of energy Q is wrongly assigned an energy in the interval $E + dE$. In Eq. (3.17) the energy dependent dispersion $\sigma_{Xe}(Q)$ is the one estimated for the Xenon detector [48]:

$$\sigma_{Xe}(Q) = \sqrt{Q/\text{KeV}} (0.579 \text{ KeV}) + 0.021 Q. \quad (3.18)$$

The observed direct detection rate is hence equal to

$$\frac{dR}{dE} = \epsilon \int_0^\infty dQ \xi(E, Q) \frac{dR}{dQ}(Q). \quad (3.19)$$

Regarding the experimental noise, we consider two contributions characterized by a different energy dependence: a flat noise component (*i.e.* constant in energy) and an exponential noise component $\propto \exp(-E/E_0)$ with $E_0 = 10$ KeV. Finally, in all the conclusions drawn below we will assume an experimental raw exposure of 1000 kg-years and a signal region ranging between 8 KeV and 75 KeV.

3.3 Bayesian forecasts

To test the capability in reconstructing the DM mass and SI cross section of a ton-scale direct detection experiment with specifications resembling the ones expected for XENON1T, we will first simulate a set of mock direct detection data and then analyze them employing a Bayesian approach supported by a Markov Chain Monte Carlo (MCMC) Likelihood scanning technique. Mock data have been simulated for each pMSSM benchmark analyzed in this work.

As already mentioned, in the context of the direct detection we need to extend the pMSSM parameter space in order to include additional parameters crucial for the computation of the differential rate (3.2). These are the parameters introduced in section 3.1 when defying the Galactic model used to compute the DM phase space density F_h within Eddington's approach. We denote by $\mathbf{p}$ the combination of the pMSSM parameters used to describe a certain benchmark and the parameters characterizing the underlying Galactic model. Similarly, by $\mathbf{p}_b$ we denote the value of $\mathbf{p}$ at the considered benchmark point. Whereas the assumed pMSSM parameters differ for the four benchmarks studied in this paper, in all cases we adopt the same benchmark Galactic model. This has parameters fixed to the mean values found in [4]. The mean local DM phase space density F_h for the Einasto, NFW and Burkert profiles can be downloaded from the online material associated with Ref. [4].

The data simulation consists in sampling a set of N recoil energies E_i , $i = 1, \dots, N$ from the probability density function $f(E, \mathbf{p}_b)$ obtained evaluating at the benchmark point $\mathbf{p}_b$ the spectral function

$$f(E, \mathbf{p}) = \frac{1}{\mu(\mathbf{p})} \left\{ \frac{dR(E, \mathbf{p})}{dE} + \frac{1}{b-a} + \frac{\exp(-E/E_0)}{E_0 [\exp(-a/E_0) - \exp(-b/E_0)]} \right\} \quad (3.20)$$

where $a = 8$ KeV, $b = 75$ KeV and $\mu(\mathbf{p}) = \int_a^b d\bar{E} dR(\bar{E}, \mathbf{p})/dE + 2$. This choice corresponds to assume two background events in the signal region during the exposure time:⁶ one from the flat noise component and one from the exponential noise component. For a given point $\mathbf{p}$ in parameter space, Eq. (3.20) represents the probability that the XENON1T experiment measures an energy (infinitesimally close to) E within the signal region (a, b) . The first term on the right hand side of this expression corresponds to the contribution to this probability from the true DM signal while the second and the third terms are associated with the flat and exponential noises respectively. The number of recoil energies N is drawn from a

⁶As expected for ton-scale direct detection experiments which aim at reducing the number of background events measured during the exposure time to ~ 1 .

Poisson distribution of mean λ_b with probability

$$\text{Poisson}(N; \lambda_b) = \frac{\lambda_b^N}{N!} \exp(-\lambda_b) \quad (3.21)$$

where $\lambda_b \equiv \mu(\mathbf{p}_b)$ is the number of events (*i.e* measured recoil energies) expected for the benchmark under analysis. In Tab. 1 we report the values of N generated in the simulations analyzed in this work.

Having explained how to generate mock data from a given benchmark point, we can now focus on their Bayesian analysis. As already mentioned the aim is to reconstruct the dark matter mass and SI cross section. We will estimate the "preferred values" for these quantities by determining their marginal posterior probability density functions (PDF). According to Bayes' theorem the posterior PDF of certain parameters $\mathbf{p}$ is proportional to the product of the Likelihood function $\mathcal{L}(\mathbf{p}, \mathbf{d})$ and the prior probability density $\pi(\mathbf{p})$:

$$\mathcal{P}(\mathbf{p}, \mathbf{d}) = \frac{\mathcal{L}(\mathbf{d}, \mathbf{p})\pi(\mathbf{p})}{\mathcal{E}(\mathbf{d})} \quad (3.22)$$

where $\mathbf{d}$ is the array of datasets used to constrain the parameter space. The Bayesian evidence $\mathcal{E}(\mathbf{d})$, being independent from $\mathbf{p}$, plays the role of a normalization constant when performing parameter inference. The marginal poster PDF of a generic function g of the parameters $\mathbf{p}$, *e.g.* σ_{SI} and M_χ in our case, is given by the expression

$$p(g|\mathbf{d}) = \int d\mathbf{p} \delta(g(\mathbf{p}) - g) \mathcal{P}(\mathbf{p}|\mathbf{d}), \quad (3.23)$$

which follows from the definition of conditional probability. MCMC scanning techniques have been then used to sample from Eqs. (3.22) and (3.23) the marginal posterior PDF of the DM mass and SI cross section.

The Likelihood function is a key ingredient of the present analysis and it has been constructed following the prescriptions given in Ref. [48]. The contribution to the Likelihood from the simulated direct detection data is given by

$$-\log \mathcal{L}_{\text{dd}}(E_i, N, \mathbf{p}) = \mu(\mathbf{p}) - N - N \log \left(\frac{\mu(\mathbf{p})}{N} \right) - \sum_{i=1}^N \log \left(\frac{f(E_i, \mathbf{p})}{f(E_i, \hat{\mathbf{p}})} \right) \quad (3.24)$$

and it has been normalized in such a way that $\log \mathcal{L}_{\text{dd}}(E_i, N, \hat{\mathbf{p}}) = 0$, where $\mu(\hat{\mathbf{p}}) = N$. This expression reflects both the information on the number of simulated recoil events N , included in the Poissonian pre-factor, and the information about their energies E_i , encoded in the spectral function (3.20). In addition to the direct detection contribution, we have included in the Likelihood a term which assigns a higher statistical weight to points in parameter space characterized by a "physical mass spectrum". This term implements in the analysis the LEP2 constraints on SUSY parameters, bounds from collider searches of Higgs decays into τ pairs (which can be interpreted as an m_A -dependent upper bound on the value of $\tan \beta$) and, finally, a lower bound of 600 GeV on the gluino mass [23]. These constraints, being expressed in terms of lower limits $\ell_r(\mathbf{d})$ on certain parameters

or theoretical predictions $\tau_r(\mathbf{p})$, generate a contribution to the Likelihood function of the following form

$$-\log \mathcal{L}_{\text{pc}}^{(r)}(\mathbf{d}, \mathbf{p}) = \begin{cases} \frac{1}{2} \left(\frac{\tau_r(\mathbf{p}) - \ell_r(\mathbf{d})}{\sigma_r(\mathbf{d})} \right)^2 & \text{if } \tau_r(\mathbf{p}) < \ell_r(\mathbf{d}) \\ 0 & \text{otherwise} \end{cases} \quad (3.25)$$

where the index r labels the constraints on the pMSSM mass spectrum previously mentioned and $\sigma_r(\mathbf{d})$ is an estimate of the error associated with the value of the lower limit $\ell_r(\mathbf{d})$. We also included in the Likelihood a term accounting for the recently discovered boson compatible with the Standard Model Higgs:

$$-\log \mathcal{L}_{\text{Higgs}}(\mathbf{d}, \mathbf{p}) = \frac{1}{2} \left(\frac{m_h(\mathbf{p}) - \hat{m}_h(\mathbf{d})}{\sigma_h(\mathbf{d})} \right)^2 \quad (3.26)$$

where $m_h(\mathbf{p})$ is the pMSSM prediction for the light CP-even Higgs boson, $\hat{m}_h(\mathbf{d}) = 126$ GeV and $\sigma_h(\mathbf{d}) = 3$ GeV. Summing up all the contributions, the final form of the Likelihood function considered in this analysis is

$$-\log \mathcal{L}(\mathbf{q}, \mathbf{p}) = -\log \mathcal{L}_{\text{dd}}(E_i, N, \mathbf{p}) - \sum_r \log \mathcal{L}_{\text{pc}}^{(r)}(\mathbf{d}, \mathbf{p}) - \log \mathcal{L}_{\text{Higgs}}(\mathbf{d}, \mathbf{p}). \quad (3.27)$$

The last information needed to complete the analysis setup concerns with the choice of the prior PDF $\pi(\mathbf{p})$. In the literature the most investigated choices correspond to flat and logarithmic priors PDF in parameter space. Regarding the pMSSM mass parameters, in the reconstruction of the DM mass and SI cross section we will consider both cases separately to assess the impact of priors on the final results. Concerning instead $\tan\beta$ and the trilinear couplings, following [49], only flat priors will be considered. To carefully sample the posterior PDF of the pMSSM parameters, we extended in the present analysis the ranges introduced in Eq. (2.1), varying squark and slepton masses between 50 GeV and 4 TeV. The ranges considered for the other parameters coincide instead with the ones of Eq. (2.1). Galactic model parameters will be treated as nuisance parameters, using the knowledge on their distributions acquired in Refs. [3, 4]: the posterior PDF found in these references for the Galactic model parameters through a compilation of various dynamical constraints will be considered in the present analysis as the prior PDF of the astrophysical parameters. As a final remark, we mention here that all the parameters controlling the hadronic matrix elements required to evaluate the DM-nucleus SI scattering cross section have been fixed at the mean values of Ref. [48]. In fact, given the high dimensionality of the pMSSM and the limited constraining power of the data, we preferred to consider in this paper as nuisance parameters only the astrophysical parameters to which we could assign prior PDF motivated by previous works. We refer the reader to Refs. [50] for an analysis assessing the role of hadronic matrix elements in the context of DM direct detection.




Figure 3: Histograms showing the distributions of the best fit DM masses derived by fitting 1000 data realizations with the theoretical expectation (3.20). The left panel refers to benchmark **xen1** and the right panel to benchmark **xen2**. The red cross represents the value of the DM mass at the benchmark point.

3.4 Statistical limitations

By reconstructing the DM mass and SI cross section we mean what follows: to sample the posterior PDF (3.22) through a MCMC algorithm⁷ and derive from it the marginal PDF of the DM mass and SI scattering cross section. We will present results in terms of $\alpha\%$ Bayesian credible regions containing therefore a fraction α of the posterior probability. Focusing on credible regions rather than on frequentist confidence intervals allows to circumvent possible issues regarding the coverage of the resulting intervals, a problem which has recently attracted a great attention in the context of the Minimal Supersymmetric Standard Model [51] and in the DM direct detection field [11].

There is however another possible source of "statistical limitations" affecting the reconstruction procedure described above. This concerns with the fact that the mock data implemented in the analysis, for instance the recoil energies E_i , represent just one single realization of the true underlying distribution of the data. If the simulated data do not provide a faithful representation of their true distribution, because of statistical fluctuations, the resulting reconstruction can be significantly biased. In Ref. [11] this effect has been observed for DM candidates covering a non negligible spectrum of masses and SI scattering cross sections. It is therefore important to check the sensitivity of our benchmark reconstructions to the particular realization of the data implemented in the analysis. Given the sophistication of the Galactic model of section 3.1 and of the particle physics parameter space considered in this work, to run several MCMC for different realizations of the data is not an efficient way of achieving this task. Therefore, to assess the dependence of our results from the use of a single data sample for each benchmark we followed a different approach based on a simple one dimensional χ^2 minimization: we first sampled for each benchmark a large number of datasets (N, E_i) as explained in the previous subsection. Then, for each

⁷In this analysis we employ a modified version of the **SuperBayes** code interfaced with **DarkSUSY** routines to compute direct detection observables.




Figure 4: As in Fig.3 but the left panel refers to benchmark **xen3** and the right panel to benchmark **xen4**.

realization of the direct detection data, we fit the resulting distribution of the N energies E_i against the theoretical expectation (3.20) varying the DM mass only and fixing the SI scattering cross section and astrophysical parameters to their benchmark values. The resulting distribution of best fit values for the DM mass will allow us to estimate the degree of sensitivity of the benchmarks studied in this paper to the specific data sample considered in the reconstruction of the DM mass and SI cross section. It is in fact the DM mass the quantity mostly affected by statistical fluctuations in the distribution of the simulated recoil energies.

The results of this analysis are shown in Fig.3 for benchmarks **xen1** and **xen2**, while Fig.4 refers to benchmarks **xen3** and **xen4**. These histograms show the distributions of the best fit DM masses derived through the fitting procedure explained above for 1000 different data realizations. In each figure the red cross represents the value of the mass at the benchmark point. In all cases we see that the distribution is not perfectly symmetric around the benchmark point: there is a general trend in overestimating the DM mass. This result is in agreement with what found in Ref. [11], where the impact of statistical limitations on the reconstruction of the DM mass and SI cross section has been studied for a broad spectrum of DM candidates. This study shows that the most sensitive DM candidates are those with high masses and low SI cross section, *i.e.* candidates associated with a relatively low number of expected direct detection signal events. This trend is confirmed by the outcome of our analysis, which shows how the benchmarks potentially more sensitive to statistical limitations are those with the highest mass, namely **xen3** and **xen4**, as clearly reported in Fig.4. The reason behind this phenomenon is that when the number of recoil events is not sufficiently large to produce a faithful description of the underlying recoil energy distribution, a small excess of events at high energy, resulting from a statistical fluctuation, is sufficient to alter the result of the fit: this excess of events at high energy will be better fitted by an artificially large value of DM mass (see also Fig.1 in Ref. [11]).

For the benchmarks studied in this paper we notice however that this effect, though




Figure 5: Two-dimensional marginal posterior PDF in the plane DM mass versus SI scattering cross section. The left panel refers to **xen1** while the right one to **xen2**. In each panel the colored bar follows the posterior PDF, arbitrarily normalized to one, from its peak down to zero. Black contours correspond to the 68% and 95% credible regions obtained implementing in the analysis flat priors for the pMSSM mass parameters, while the green contours correspond to the analogous quantities derived from log-priors. The white dot and the blue cross represent respectively the posterior mean (from log-priors) and the benchmark point.

present, it does not influence dramatically the best fit distributions of the DM mass. Since therefore we do not observe any strong pathology related to the selected benchmarks, we continue our analysis focusing on their mass and SI cross section reconstruction at XENON1T.

3.5 Mass and cross-section reconstruction

The procedure to reconstruct the DM mass and SI cross section introduced in the previous subsections has been applied to the benchmarks **xen1**, **xen2**, **xen3** and **xen4**. For each benchmark we focused on a single data realization considering both flat and log-priors for the pMSSM mass parameters. In Fig. 5 and 6 we show for these benchmarks the two-dimensional marginal posterior PDF in the plane DM mass versus SI scattering cross section. In each panel the colored bar follows the two-dimensional posterior PDF which has been conveniently normalized to one. Black contours correspond to the 68% and 95% credible regions obtained implementing in the analysis flat priors for the pMSSM mass parameters, while the green contours correspond to the analogous quantities derived from log-priors. Finally, the white dot and the blue cross represent respectively the posterior mean (from log-priors) and the benchmark point.

The plots confirm the general tendency in overestimating the DM mass described in the previous subsection, here evidenced by the high mass tail featured by the contours and partially by the mismatch between the posterior means and the values of the mass and




Figure 6: As in Fig.5 but the left panel refers to benchmark **xen3** and the right panel to benchmark **xen4**.

SI cross section of the original benchmark. On the other hand we observe a significant improvement of the reconstruction performance when adopting log-priors in the MCMC scanning procedure. Indeed, this choice tends to “compensate” possible limitations in the DM mass reconstruction, due to statistical fluctuations in the distribution of the simulated data, by giving more statistical weight to low values of the parameters [52] (and therefore to the DM mass), thus partially reducing the high mass tail of the marginal posterior PDF. Because of this compensation mechanism we argue that log-priors provide in this context a more faithful description of the actual potential of upcoming 1-ton DD experiments and we will therefore employ results obtained with this choice when translating our DD analysis into a forecast for the LHC.

For a given benchmark, in addition to this effect, the quality of the reconstruction is also affected by the value of the DM mass itself. Indeed, the differential recoil rate, for DM masses largely exceeding the one of the nuclei in the target material, depends on the DM parameters only through their ratio σ_{SI}/M_χ . Clearly, this degeneracy produces credible regions spanning over “erroneously” large portions of the plane (M_χ, σ_{SI}) . In the present analysis the benchmarks most largely affected by this degeneracy are benchmarks **xen3** and **xen4**, as also shown in Fig. 6.

In the next section we will translate the preferred regions in the plane DM mass versus SI cross section determined here into a prediction for certain classes of signals at the LHC. Since most of the relevant SUSY production cross sections are very steep functions of the pMSSM mass spectrum, and therefore of the DM mass, also the LHC analysis described in the following will be indirectly affected by the statistical limitations producing large uncertainties in the DM mass reconstruction. Large exposures are therefore crucial to enhance the effectiveness of the strategy illustrated here to combine the direct detection technique with DM searches at the LHC.

In summary, a ton-scale experiment resembling the XENON1T specifications will be able to achieve a robust determination of mass and SI cross section for DM candidates similar to the one associated with the benchmark point **xen1**. In fact, in light of our simulations, **xen1** clearly emerges as the most promising benchmark considered in this work. This result follows from the large number of recoil events in the simulation of **xen1** (170) and from the fact that light candidates are in this context simpler to identify with the direct detection technique, having a steeper energy spectrum which minimizes the previously mentioned statistical limitations. The benchmark points **xen2**, **xen3** and **xen4** feature instead a less efficient reconstruction but still in moderate agreement with the true values of the DM parameters related to the original benchmark. These points are in fact more sensitive to the already mentioned statistical limitations and to the mass/cross section degeneracy, typical of heavy DM candidates.

4 Collider analysis

Having derived for each benchmark the accuracy within which the relevant DM properties are reconstructed by a 1-ton DD experiment resembling XENON1T, we now focus on how to translate this information into a prediction for certain LHC observables. Our analysis consists in two steps: first we will reprocess the Markov chains produced in the DD analysis applying to the models in the chains additional constraints from the relic density, limits on the muon anomalous magnetic moment as well as bounds from B-physics processes. Then, when possible, we will determine the production processes most likely associated to a subsample of points in the chains passing these additional constraints and study the corresponding LHC response in its 7 TeV center of mass energy configuration. This procedure is the essence of the method proposed in this paper to correlate the DD technique with DM searches at the LHC.

To rigorously assess the impact on the DD analysis of section 3 of additional observables one should run new MCMC scans with an extended Likelihood properly incorporating the new constraints. Although this is by now technically achievable in the most simple SUSY realizations (see [52, 53] for some examples) this procedure would be computationally troublesome to implement in the present analysis because of the need of repeated lengthy computations of quantities like the DM relic density, especially in the context of a high dimensional MSSM parameter space, as the one considered in this work, and when studying points in parameter space achieving the correct relic density via co-annihilation processes. For these reasons we have employed the less computationally demanding procedure of applying a posteriori the limits from the relic density and flavor processes to the Markov chains generated in the analysis of section 3. More precisely, for each benchmark we have computed the observables just mentioned only for the PDF samples enclosed in the 95% log-prior contours reported in Figs. (5) and (6). The limits implemented for δa_μ , $b \rightarrow s\gamma$ and $B_s \rightarrow \mu^+\mu^-$ are the same described in section 2. Regarding the relic density, instead, we have just required that it does not exceed the cosmological limit. This is motivated by the fact that too strict requirements may excessively affect set of points not informative on

Search channel	Main processes	References
3 leptons + E_T^{miss}	$\chi_2^0 \chi_1^\pm \rightarrow WZ + E_T^{\text{miss}}$	[62]
2 leptons + E_T^{miss}	$\chi_1^\pm \chi_2^0, \chi_1^\pm \chi_1^\mp, \tilde{l}^\pm \tilde{l}^\mp$	[63]
3 b-jets + E_T^{miss}	$\tilde{g}\tilde{g}$	[64]
≥ 1 b-jet + ≥ 4 jets + 1 lepton + E_T^{miss}	$\tilde{t}_1 \tilde{t}_1^*$	[65]
1-2 b-jets + 4-5 jets + E_T^{miss}	$\tilde{t}_1 \tilde{t}_1^*$	[66]
monojet + E_T^{miss}	$\tilde{t}_1 \tilde{t}_1^*$	[67]
2 b-jets + E_T^{miss}	$\tilde{b}_1 \tilde{b}_1^*$	[68]
2 jets + Missing Energy	$\tilde{q}\tilde{q}^*$	[69, 70]

Table 2: Summary table of the SUSY searches employed in our collider analysis.

this observable⁸.

As already mentioned, the additional constraints imposed on the model parameters restrict the 95% credible regions determined in section 3 and produce the subregions on which our collider analysis has been mostly focused. We have then identified the production processes most likely associated to the SUSY models contained within these subregions and employed accordingly the analysis strategies applied by the ATLAS [59] collaboration to a few points sampled from them. Indeed, for each of these points we have simulated the signal possibly detectable by the ATLAS experiment and obtained the number of expected signal events for several combinations of final states, as summarized in tab. 2, at the luminosities relative to the current available data. From the comparison with the current limits we have determined whether the various configurations are already severely constrained or will be probed by the very next upgrades in luminosity.

The signal events have been generated through the numerical package Madgraph [60] combined with the detector simulator PGS normalizing the production cross-section at the NLO through the package PROSPINO [61]. The simulations have been validated by generating the main backgrounds for the considered search channels and cross-checking our results with the number of background events inferred by the data. The number of events obtained by the simulations has resulted compatible with expectations.

4.1 Analysis of xen1

We start our analysis with the lightest DM mass scenario, namely **xen1**. Fig. 7, obtained selecting points in our Markov chains passing flavor and relic density constraints, evidences that values of the DM relic density compatible with current limits can be achieved in two setups only: the first is represented by the two narrow regions lying at values of the DM mass of around 45-60 GeV, thus coinciding with the original benchmark scenario. In this regions a bino-like DM candidate is driven to the correct relic density by resonance enhanced

⁸ Alternatively one could also consider deviations from the standard WIMP paradigm, as enforced by non thermal production [54–56] or alternative cosmologies [57, 58]. This would require however a different treatment of the local DM density [53]. However, SUSY parameters are less directly correlated in these scenarios.




Figure 7: Points of the Markov chains associated to the benchmark **xen1** passing flavor and relic density constraints in the plane $(M_\chi, \Omega h^2)$ following a color pattern related to the DM higgsino fraction.

interactions mediated either by the light CP-even Higgs boson or, for slightly lower mass values, by the Z boson. In the second setup, instead, occurring at higher values of the DM mass and corresponding to the yellow region in the plot, DM is mainly a pure higgsino characterized by very low values of the relic density and by a value of the mass larger than 100 GeV as required by the LEP bound.⁹

Notably, there is a clear correspondence between the second setup, represented by the yellow band in Fig. 7, and the high mass tail of the two-dimensional marginal posterior PDF in the plane DM mass versus SI scattering cross section shown in Fig. 5. As argued in section 3.4, this tail arises as a consequence of statistical fluctuations in the recoil energy spectrum affecting the accuracy of the DM mass reconstruction. We therefore conclude that the higgsino DM setup can be effectively disentangled from the bino DM case (corresponding to the original benchmark) by improving the performance of the DD based parameter inference, for instance, by increasing the experimental exposure. In view of this consideration we have limited our collider analysis to a subset of models sampled from the PDF of section 3 and corresponding to gaugino DM.

For light DM candidates, as the one considered in this section, the most relevant parameters entering the SI cross section and relic density calculation are the DM mass itself and the value of the μ parameter. This implies that, from the collider point of view, the most interesting way to probe this scenario consists in the search for direct gaugino production,

⁹Fig. 7 also reports a very small amount of points at gaugino DM in the mass interval of around 70-300 GeV. For these points a correct value for the relic density is guaranteed by either a low mass slepton or by a CP-odd Higgs. In the first case, however, the slepton should lie very close to the LEP limit and it is thus statistically disfavored by our Monte Carlo scanning procedure, while in the second low values of m_A are disfavored by $Br(B_s \rightarrow \mu^+ \mu^-)$.

benchmark	M_2	μ	$\sigma_{\text{vis}}[fb]$ SR1	$\sigma_{\text{vis}}[fb]$ SR2
xen1-1	498	283	0.1	0.3
xen1-2	215	362	0.3	0.6
xen1-3	212	240	1.7	3.3
xen1-4	138	263	1.6	0.1
xen1-5	266	240	0.5	1.2

Table 3: Values of the non-SM visible cross section associated to 3 leptons + E_T^{miss} search channel for 5 sample points relative to the light DM scenario depicted by the benchmark **xen1**. The results rely on two signal regions associated, respectively, to the decay of the gauginos into off-shell and on-shell gauge bosons. The experimental limits are $\sigma_{\text{vis}}^{\text{obs}} = 3 \text{ fb}$ and $\sigma_{\text{vis}}^{\text{obs}} = 2 \text{ fb}$.

namely the associated production of χ_2^0 and $\chi_1^\pm$. Assuming that there are no intermediate sleptons between these two states, they can decay either into $W^\pm Z$, with the bosons which may be also off-shell, or in the pair Wh , in addition to missing energy. Our study has been focused on the first possibility employing the ATLAS dedicated search through events with 3 leptons in the final state [62]. A strategy for the detection of the Wh channel has been instead proposed in [71] but it requires much higher luminosities for being detected compared to the previous channel.

Processes involving the production of colored particles do not play a relevant role, at least up to moderate values of M_2 , since the current collider limits imply heavy masses such that the relative production rates can be regarded as subdominant. Notice that this, in the pMSSM, is strictly valid for the gluino and the squarks of the first two generations; third family squarks can feature much less stringent limits¹⁰. This is likely not the case for such low values of the DM mass. Indeed sbottom masses are excluded up to around 390 GeV at 2.05 fb^{-1} for DM masses less than 60 GeV [75] and preliminary results from the 4.7 fb^{-1} update [68] move the lower limit up to around 500 GeV. Regarding the stop (see, for reference, [65, 66, 76]), some values in the low mass range are still not excluded (the reason is that the stop features several different decay channels and then the study of very different signals, each one probing a limited range of masses, is required. More details will be provided in the following sections). However, for the values of the SI cross section emerging in the present scenario, squark mediated interactions are not expected to play an important role, then it is more interesting to focus on collider searches probing instead the relevant parameters, as for instance the previously mentioned 3 lepton final state.

Tab. 3 reports the results relative to 5 models, named **xen1-1**, ..., **xen1-5**, sampled from the PDF relative to **xen1** and passing relic density and flavor constraints. The table reports, for each model, the values of M_2 and μ which determine the characteristics of the produced neutralino and chargino, together with the visible cross-sections for non-SM

¹⁰A part from the cases discussed in the next sections, in which it is established a correlation with the relic density requirement, we exclude as well the possibility of very compressed spectra (see e.g. [72–74] and references therein for some studies).




Figure 8: Missing energy distribution for the events (signal+background) including three leptons in the final state expected for the models **xen1-1**, ..., **xen1-5** sampled from the chains relative to benchmark **xen1**. This result has been obtained in a run of the LHC with an energy in the center of mass of 7 TeV and a luminosity of 4.7 fb^{-1} .

processes, i.e. the product $\sigma \epsilon A$ of the theoretical production cross-section σ , the detector efficiency ϵ and the acceptance A which accounts for the reduction of the signal strength due to the analysis cuts. DD combined with relic density and flavor requirements has a good constraining power for the μ parameter, which we have found to vary on a moderately restricted range with a lower bound of around 200 GeV, as expected from the functional form of the SI cross section, while leaves substantially M_2 free to vary. The results in the table shows that the WZ channel can efficiently probe models with moderate values of M_2 with some regions already excluded and some other which can be detected in the near future with an $O(1)$ increase of the luminosity. At high values of M_2 , instead, the chargino and the second neutralino increase their higgsino fraction favoring the decay into Wh and requiring therefore a different search strategy compared to the one considered here involving three leptons in the final state.

Fig. 8 shows the missing energy distribution corresponding to events (signal+ background) including three leptons in the final state simulated from the models **xen1-1**, ..., **xen1-5** in a run of the LHC with an energy in the center of mass of 7 TeV and a luminosity of 4.7 fb^{-1} . In summary, the benchmark scenario **xen1** features a clearly identifiable signature in the three lepton final state emerging as a genuine forecast of our DD analysis which in this case can be effectively translated into the results of Fig. 8 and Tab. 3.

4.2 Analysis of xen2

The benchmark point **xen2**, similarly to **xen1**, has been designed to study MSSM realizations featuring a light EW interacting sector (see e.g. [77] for a similar approach) evading current constraints which would instead apply to models including colored sectors lying in




Figure 9: Points of the Markov chains associated to the benchmark **xen2** passing flavor and relic density constraints in the plane $(M_\chi, \Omega h^2)$ following a color pattern related to the DM higgsino fraction.

the multi-TeV region. As for the case of **xen1**, in a preliminary investigation of the benchmark **xen2**, we selected from our Markov chains the models passing flavor and relic density constraints. These points are shown in Fig. 9 where again the yellow band corresponds to higgsino DM configurations. Contrary to the case of **xen1**, however, in the present scenario the higgsino DM models do not only populate the tail of the marginal posterior PDF in the plane DM mass versus SI scattering cross section: comparing in fact Fig. 5 with Fig. 9 one can clearly observe that there are regions in the plane (M_χ, σ_{SI}) with a significant statistical weight corresponding to DM masses above 100 GeV and therefore representing a large fraction of points in the yellow band of Fig. 9. This is obviously related to the fact that the LEP bound implemented in the Likelihood of our MCMC scans is only effective for exotic states lighter than 100 GeV, while the DM candidate associated with **xen2** has a mass of approximately 155 GeV.

For these reasons the outcome of our DD based analysis cannot resolve alone the degeneracy among pure gaugino and pure higgsino DM configurations (see Fig. 9). A possible strategy to alleviate this problem could be to encode in the Likelihood the requirement of a DM relic density lying within $2-3\sigma$ from the cosmological limit; this requirement cannot be fulfilled by higgsinos with relatively low values of the mass, as required by this benchmark scenario. A potentially interesting alternative is, on the other hand, adding information coming from indirect detection. Indeed bino-like and higgsino DM feature different signatures from the point of view of indirect detection. In particular, current experimental facilities have reached enough sensitivity to probe light higgsino DM scenarios with already strong constraints coming from antiproton data [78] and new stronger bounds eventually arising in case that the claim of detection of a gamma-ray line in the Fermi-LAT data [79]

benchmark	m_χ	M_2	μ	$m_{\tilde{l}}$	$m_{\tilde{\tau}}$	$\sigma_{\text{vis}}[fb] SR1$	$\sigma_{\text{vis}}[fb] SR2$
xen2-1	132	348	356	256	139	0.13	0.09
xen2-2	155	462	940	180	1100	1.24	0.20
xen2-3	175	187	554	370	2500	0.03	0.02
xen2-4	140	318	388	160	2300	0.18	0.11
xen2-5	160	3431	620	2745	182	0.001	0.003

Table 4: Values of the non-SM visible cross sections associated to the 3 leptons + E_T^{miss} search channel for 5 sample points relative to the DM scenario depicted by the benchmark **xen2**. The results rely on three of the signal regions considered in Ref. [62]. The experimental limits are, respectively, $\sigma_{\text{vis}}^{\text{obs}} = 3 \text{ fb}$ and $\sigma_{\text{vis}}^{\text{obs}} = 2 \text{ fb}$.

benchmark	$\sigma_{\text{vis}}[fb] SR_{MT2}$	$\sigma_{\text{vis}}[fb] SR_{OS}$	$\sigma_{\text{vis}}[fb] SR_{SS}$
xen2-1	1.63	1.67	0.07
xen2-2	0.10	0.16	0.07
xen2-3	0.39	0.57	0.04
xen2-4	0.14	0.21	0.02
xen2-5	0.01	0.02	0.01

Table 5: Values of the non-SM visible cross sections associated to the 2 leptons + E_T^{miss} search channel for 5 sample points relative to the DM scenario depicted by the benchmark **xen2**. The results rely on three of the signal regions considered in Ref. [63]. The experimental limits are, respectively, $\sigma_{\text{vis}}^{\text{obs}} = 2 \text{ fb}$, $\sigma_{\text{vis}}^{\text{obs}} = 11.4 \text{ fb}$ and $\sigma_{\text{vis}}^{\text{obs}} = 2 \text{ fb}$.

will be confirmed (for more details see e.g. [80]).

Leaving these possibilities for future works, we will focus in this section on the scenario corresponding to gaugino DM while a discussion on possible peculiar signatures of higgsino DM at the LHC is postponed to the next section. Moreover, we will restrict in the present collider analysis to the DM mass range 135-175 GeV, corresponding to a region of high posterior PDF in the plane DM mass versus scattering cross section able to capture the main features of the present scenario. Within this mass range there are several different possibilities to achieve a DM relic density compatible with the cosmological limit. The main ones are the coannihilation with a slepton and resonances occurring in annihilations mediated by the CP-odd Higgs, with the latter challenged by the constraints from $B_s \rightarrow \mu^+ \mu^-$. In addition to these, we have also found configurations in which the parameters M_1 and M_2 are very close to each other, featuring the already mentioned setup dubbed “well tempered neutralino”: in this case the correct relic density is achieved by the interplay of a non negligible wino fraction of the DM and its coannihilations with the lightest chargino and the second neutralino.

This variety of scenarios consequently requires a combination of different collider strategies in order to disentangle the larger number of possible signals. Analogously to the previous




Figure 10: Missing energy distribution for the events (signal+background) including two leptons in the final state expected for the models **xen2-1**, ..., **xen2-5** sampled from the chains relative to benchmark **xen2**. This result has been obtained in a run of the LHC with an energy in the center of mass of 7 TeV and a luminosity of 4.7 fb^{-1} . The left panel refers to the signal region dubbed MT2 while the right panel corresponds to the signal region OS defined in Ref. [63].

benchmark we still stick on the direct production of EW interacting particles. In addition to three-lepton signals, however, we have also employed detection strategies of events featuring two leptons in the final state [63] which improve the sensitivity of the previous search in the case of the presence of sleptons in the decay chain of $\chi_1^\pm$ and are, moreover, sensitive to direct slepton production as well as other gaugino production processes like $\chi_1^\pm \chi_1^\mp$ which can be potentially relevant.

The outcome of our analysis, performed on 5 representative points (denoted by **xen2-1**, ..., **xen2-5**) sampled from the posterior PDF of **xen2** and reported in Tabs. (4) and (5), shows that the collider prospects are mainly influenced by the value of M_2 and by the slepton mass scales while the μ parameter, partially fixed by DD, is important in determining the composition of $\chi_1^\pm$ and χ_2^0 . It is also evident that the analysis employed are not sensitive to low mass splittings among the relevant particles. Results of a LHC run simulated from these points focused on final states including two and three leptons are shown in Fig. 10.

We conclude this subsection commenting again on the possible presence of light third family squarks. The higher value of the DM mass (compared to **xen1**) weaken sensitively the current limits, then in principle they could account for relevant signals. However in the setup we are considering they do not impact the DM relic density since they cannot be enough light to give rise to coannihilations (concerning this claim, see also the outcome of the next subsection). For these reasons we will not consider this possibility since our principal aim is to investigate MSSM realizations which enforce correlations among the different experimental observables. Moreover, we notice that also in the present benchmark scenario we are exploring regions of the parameter space where the SI cross section is




Figure 11: Points of the Markov chain associated to the benchmark **xen3**, selected according to the criteria depicted in the text and passing flavor and relic density constraints. In each panel the points are reported in the plane $(m_\chi, \Omega h^2)$. In the upper left panel the points are colored according to the corresponding value of the DM higgsino fraction. In the upper right panel instead the color pattern is determined by the quantity $\Delta_A = (m_A - m_\chi)/m_A$. The bottom left and bottom right panels are obtained in analogous manner but the quantities discriminated by the colors are respectively $\Delta_b = (m_{\tilde{b}_1} - m_\chi)/m_{\tilde{b}_1}$ and $\Delta_t = (m_{\tilde{t}_1} - m_\chi)/m_{\tilde{t}_1}$.

dominated by the parameters of the Higgs sector. In absence of further constraints (for example coming from the relic density), the scan on the mass of the third family squarks (performed in the DD analysis) is uniquely influenced by the bound on the Higgs mass implemented in the Likelihood function which tends to push them towards high values (see also the next section for further details on this aspect).

4.3 Analysis of xen3

This benchmark is designed to probe coannihilation scenarios with third family squarks very close to the current experimental limits relative to their direct collider production. As already mentioned, the contribution to the SI cross section of the squark mediated

benchmark	m_χ	$m_{\tilde{t}_1}$	$m_{\tilde{b}_1}$	
xen3-1	250	524	270	$\tilde{b}_1 \rightarrow b\chi_1^0, \tilde{t}_1 \rightarrow t\chi_1^0$
xen3-2	230	256	442	$\tilde{b}_1 \rightarrow b\chi_1^0, \tilde{t}_1 W^- \tilde{t}_1 \rightarrow c\chi_1^0$
xen3-3	219	249	301	$\tilde{b}_1 \rightarrow b\chi_1^0, \tilde{t}_1 \rightarrow c\chi_1^0$
xen3-4	218	605	234	$\tilde{b}_1 \rightarrow b\chi_1^0, \tilde{t}_1 \rightarrow t\chi_1^0$
xen3-5	198	356	217	$\tilde{b}_1 \rightarrow b\chi_1^0, \tilde{t}_1 \rightarrow \tilde{b}_1 W^+$
xen3-6	150	246	164	$\tilde{b}_1 \rightarrow b\chi_1^0, \tilde{t}_1 \rightarrow \tilde{b}_1 W^+$
xen3-7	182	540	196	$\tilde{b}_1 \rightarrow b\chi_1^0, \tilde{t}_1 \rightarrow t\chi_1^0$
xen3-8	165	511	175	$\tilde{b}_1 \rightarrow b\chi_1^0, \tilde{t}_1 \rightarrow t\chi_1^0$
xen3-9	172	343	181	$\tilde{b}_1 \rightarrow b\chi_1^0, \tilde{t}_1 \rightarrow \tilde{b}_1 W^+$
xen3-10	220	232	380	$\tilde{b}_1 \rightarrow b\chi_1^0, \tilde{t}_1 \rightarrow c\chi_1^0$

Table 6: Summary table of the points relative to the scenario **xen3** used for the collider analysis. The last column reports the main decay channels of the lightest stop and sbottom in the models considered.

interactions is enhanced for low DM-squark mass splittings and it dominates the cross-section itself in the limit of very low higgsino fraction. At the same time these low mass splittings enforce coannihilations driving the DM relic density towards its correct value and imply very simple and definite decay chains at collider.

As for the other benchmark points, we start the analysis of **xen3** reprocessing the Markov chains produced by the DD simulation applying to the sampled models the current bounds from the DM relic density and flavor physics. The points passing these bounds are shown in Fig. 11, where the color patterns follow the higgsino fraction (top left panel), $(m_A - m_\chi)/m_A$ (top right panel), Δ_b (bottom left panel) and, finally Δ_t (bottom right panel). The outcome of this analysis shows that extremely pure higgsino DM configurations are favored by the interplay of the DD simulation and the additional constraints subsequently imposed on the resulting Markov chains. Moreover, bino-like DM configurations passing these additional constraints (and corresponding to the original benchmark scenario) achieve the correct relic density mainly through the Higgs funnel mechanism while coannihilations are disfavored by the requirements on the Higgs mass (a sizable fraction of points in the chains is also excluded by the bounds from $b \rightarrow s\gamma$ and $B_s \rightarrow \mu^+\mu^-$). This outcome confirms the difficulty, already mentioned in section 2, of realizing configurations in which the SI cross section is dominated by squark-mediated interactions and thus points again towards the need of combining the information coming from DD with the fit of some other observable, like for instance the relic density, which selects more efficiently the coannihilation scenario.

The different scenarios emerging from Fig. 11 and corresponding to bino-like DM and higgsino-like DM candidates thermally produced in various ways (not only through coannihilations with light squarks) have different collider signatures. Since therefore in light of our DD simulation we can not associate to the present benchmark scenario a single clean

benchmark	SR1	SR2	SR3	SR4
$\sigma_{\text{vis}}^{\text{obs}}[\text{pb}]$	1.63	0.13	0.026	0.006
xen3-1	0.2	0.04	0.005	0.0008
xen3-2	0.25	0.04	0.005	0.001
xen3-3	0.3	0.05	0.007	0.001
xen3-4	0.4	0.07	0.010	0.0015
xen3-5	0.5	0.08	0.010	0.001
xen3-6	1.59	0.19	0.025	0.002
xen3-7	0.98	0.13	0.019	0.002
xen3-8	1.66	0.23	0.035	0.007
xen3-9	1.37	0.18	0.027	0.004
xen3-10	0.51	0.08	0.011	0.002

Table 7: Values of the non-SM visible cross sections of the points used for the collider analysis with respect to the 4 signal regions defined in Ref. [67]. This analysis is aimed at exploring processes leading to sensitively soft jets and then is particularly suitable to probe coannihilation scenarios.

prediction for the LHC, as for instance in the case of **xen1**, we will consider in the following different cases separately, arguing that, in spite of the already mentioned difficulties, the coannihilation setup involving a bino-like DM candidate and a stop or a sbottom NLSP has different collider prospects compared to, for instance, configurations corresponding to higgsino-like DM candidates. Analogously to the case of **xen2**, we will restrict to regions of the (M_χ, σ_{SI}) plane exhibiting a high statistical weight and capturing all the features relevant for the present analysis, namely a region corresponding to DM masses lower than 250 GeV.

We now focus on the bino-like configurations emerging from our Monte Carlo scans. Given the DM mass range relevant for the present analysis, production processes involving gluinos and first generation squarks are still severely constrained. If stops and sbottoms lie in the TeV range, like in the CP-odd resonance scenario, the dominant process in this case is again gaugino production with typical cross section of only few fb and thus very difficult to detect in the next future. For this reason we will not further consider this possibility in what follows. On the contrary the coannihilation scenario involving sbottom and squarks NLSP features sizable production rates for these particles. A coannihilating sbottom NLSP can only decay through the channel $\tilde{b} \rightarrow b\chi_1^0$ while in the case of a stop NLSP the dominant decay channel is often $t \rightarrow c\chi_1^0$, arising at one-loop, while the other possible decay is $\tilde{t} \rightarrow b\bar{f}f'\chi_1^0$, with f and f' being two light fermions. These decays, however, evade most of the current SUSY searches since their products are too soft to pass the analysis cuts on the p_T , or even the requirements for correct reconstruction of the final states. Such configurations thus require search strategies optimized for very low mass splittings or alternatively can be probed using initial state radiation. In this work we have

benchmark	SRAtight	SRAmedium	SRA'
$\sigma_{\text{vis}}^{\text{obs}} [\text{fb}]$	0.66	5.1	6.0
xen3-1	0	0.1	0.2
xen3-2	0	0.1	0.1
xen3-3	0.04	1.2	2.2
xen3-4	0	1.2	1.7
xen3-5	0.2	1.2	2.2
xen3-6	0.2	2.8	4.6
xen3-7	0.07	1.8	2.9
xen3-8	0.2	3.6	5.7
xen3-9	0.08	2.1	4.1
xen3-10	0.09	1.1	2.1

Table 8: Values of the non-SM visible cross sections of the points used for the collider analysis with respect to the 3 signal regions at 2-jet minimal multiplicity defined in Ref. [69]. This analysis is aimed at exploring decay chains with low mass splittings.

focused on the case of the initial radiation of a jet employing the dedicated ATLAS search of Ref. [81]. Good prospects of probing scenarios with very low mass splitting have been obtained as well from the study of monophoton events in Ref. [74].

To this purpose we have sampled from the PDF obtained from our DD analysis a few representative models (listed in Tab. 6) and generated from them LHC final states involving pairs of stop and sbottom squark with an additional jet. The results, reported in Tab. 7, show that this coannihilation scenario is rather efficiently probed, with some points already excluded, especially at low masses while we have a significant decrease in the performance at higher values since the stop/sbottom production cross sections are rather steeply decreasing functions of the masses of these particles. These results further stress the importance of increasing the performance in reconstructing the DM mass (in this case related by coannihilation requirements to the squark ones) since even slightly different determinations drastically alter the collider detection prospects.

The results of monojet searches can be interpreted in a large variety of models including some non-SUSY extension of the SM, as for instance the one considered in Ref. [81]. In addition to the monojet analysis, we have then employed a search based on two jets in the final state (see also Ref. [82]) featuring a signal region referred as A' [69] specifically designed for being sensitive to low particle mass splittings¹¹. As evident from Tab. 8, this kind of search has shown a good sensitivity for some of the sample points although it turned out to be in general less effective than the monojet analysis.

We conclude this section focusing on the collider signatures of the higgsino DM scenarios

¹¹A preliminar analysis updated to the 2012 data set with 8 TeV of center of mass energy is already available [70]. However a slightly different set of cuts have been employed. The result here shown are related to the 7 TeV data set, no sensitive improvement have been found once performing the update in luminosity and center of mass energy.

benchmark	$m_{\chi_1^0}$	$m_{\chi_1^\pm}$
HD1	134	138
HD2	145	148
HD3	169	172
HD4	175	176
HD5	194	196
HD6	211	213
HD7	234	236
HD8	247	249

Table 9: Mass of the lightest neutralino and chargino for the additional points at higgsino DM for which no additional requirement on the rest of the spectrum is imposed.

benchmark	SR1	SR2	SR3	SR4
$\sigma_{\text{vis}}^{\text{obs}}[\text{pb}]$	1.63	0.13	0.026	0.006
HD1	0.08	0.01	0.002	0.0004
HD2	0.07	0.01	0.002	0.0004
HD3	0.04	0.008	0.001	0.0002
HD4	0.04	0.008	0.001	0.0003
HD5	0.03	0.006	0.001	0.0002
HD6	0.02	0.005	0.001	0.0002
HD7	0.01	0.004	0.0008	0.0001
HD8	0.01	0.003	0.0007	0.0001

Table 10: Values of the non-SM visible cross section of the points reported in tab. (9) relative to the 4 signal regions designed for the collider search described in [67].

emerging from our Monte Carlo scans. The main peculiar feature of these SUSY spectra is the presence of a chargino and a second neutralino almost degenerate with the LSP, with the masses of all the three particles controlled by the μ parameter. The correct DM relic density is guaranteed by the higgsino nature of the DM candidate, so no other requirement is imposed on the rest of the spectrum. If this is the case, collider phenomenology is characterized again by neutralino/chargino production (namely the production of $\chi_1^\pm \chi_2^0$, but also $\chi_1^\pm \chi_1^0$ and $\chi_1^0 \chi_1^0$) as follows by the mass degeneracy of the three lightest higgsinos, thus featuring a possible signal into monojets. We have accordingly simulated the production of a pair of lightest neutralino/charginos with one additional jets for a sample drawn from our chains of eight points (see Tab. 9) with mass as low as 134 GeV in order to take into account the outcome of the previous section as well. The values of the visible cross sections are however significantly lower (see Tab. 10) with respect to the previously discussed coannihilation scenario which is then better probed by monojet searches.




Figure 12: Points in the Markov chain associated to the benchmark **xen4** passing flavor and relic density constraints. The points in the two plots follow color patterns determined, respectively, by higgsino fraction and Δ_A . By comparing the two panels one notices that a large fraction of points at gaugino DM clusters around the value $\Delta_A \simeq 0.5$.

4.4 Analysis of xen4

The last benchmark point considered in the analysis has been also designed, similarly to **xen3**, to investigate a scenario where DM is produced through coannihilations involving squarks of the third family. In this case, however, the DM mass is significantly higher than in the case of the previous benchmark. Fig. 12 clearly shows that no coannihilation configurations emerged from the interplay of the DD simulations and the application of the relic density and flavor bounds, being the PDF samples strongly dominated by higgsino DM candidates and, to some extent, by CP-odd Higgs funnel configurations. In view of this outcome, we regard the reconstruction of this point as unsuccessful and thus we will not pursue any collider analysis for this scenario.

5 Conclusions

In this paper we have explored a new strategy to combine direct and collider dark matter searches: employing the potential of the upcoming generation of 1-ton direct detection experiments, we have studied the possibility of translating the information encoded in an hypothetically discovered direct detection signal into certain classes of expected signals at the LHC.

The analysis has been performed in the context of the so-called phenomenological Minimal Supersymmetric Standard Model where a large number of possible supersymmetric configurations is conveniently described by a limited number of parameters. We focused on four benchmark points representative of WIMP dark matter candidates which should be soon probed by the upcoming generation of 1-ton direct detection experiments and corresponding to different dark matter production scenarios. In all cases a clear correla-

tion pattern between direct detection and LHC observables is enforced by the underlying mechanism accounting for the dark matter relic density.

For each benchmark, we tested the capability of a 1-ton direct detection experiment resembling the XENON1T experiment in reconstructing the dark matter mass and spin-independent cross section within the framework of a Bayesian analysis supported by a Markov Chain Monte Carlo scanning technique. In our analysis we carefully addressed the impact on the reconstructions of the astrophysical uncertainties entering the computation of the differential count rates, of possible statistical limitations related to the finite number of measured recoil energies as well as of the choice of the priors for the model parameters.

We derived through this procedure the accuracy within which the dark matter mass and scattering cross section are reconstructed by the Xenon experiment in its 1-ton configuration. We then focused on the regions of the plane dark matter mass versus scattering cross section enclosing the 95% of the posterior probability. Within these regions we selected a subset of models by reprocessing our Markov chains introducing additional constraints from the relic density, the anomalous magnetic moment of the muon and B-physics processes. For some of these models we simulated the ATLAS detector response in its configuration with 7 TeV center of mass energy and carefully studied the prospects of detecting certain classes of final states for which dedicated search strategies have been developed by the ATLAS collaboration.

The procedure proposed here defines a method to efficiently exploit the upcoming generations of 1-ton direct detection experiments to forecast valuable predictions for the LHC and it constitutes the main achievement of this paper. The results obtained within this framework are particularly promising for the benchmark **xen1** which is representative of a light bino-like neutralino produced via resonant annihilations. In this case a definite prediction for a LHC final state including three leptons and missing energy can be elaborated starting from the informations derived from the simulated 1-ton direct detection experiment. The other benchmarks, instead, are more sensitive to the direct detection statistical limitations discussed in the text and affecting the accuracy of the dark matter mass reconstruction. Larger exposures in the direct detection (here we considered 1000 kg-years) and eventually more informative Likelihoods can however further enhance the effectiveness of our approach and allow to draw interesting conclusions also in the case of challenging dark matter candidates as for instance **xen4**.

In summary, we have proposed a strategy to systematically translate an hypothetical direct detection signal into a prediction for the LHC. This method has the potential to significantly strengthen the complementarity between the direct detection technique and dark matter searches at collider and will be further investigated in future works.

Acknowledgments

It is a pleasure to thank Antonio Masiero for his collaboration during the early stages of the project and useful discussions on the main idea behind this work. We also thank Umberto de Sanctis, Alberto Tonero, Michele Pinamonti, Andrea De Simone and Florian Bonnet for fruitful conversations. These results have been partially obtained making use of the bwGRiD

Cluster (<http://www.bw-grid.de>), member of the German D-Grid initiative, funded by the Ministry for Education and Research (Bundesministerium für Bildung und Forschung) and the Ministry for Science, Research and Arts Baden-Württemberg (Ministerium für Wissenschaft, Forschung und Kunst Baden-Württemberg). G.A and P.U. acknowledges partial support from the European Union FP7 ITN-INVISIBLES (Marie Curie Actions, PITN-GA-2011- 289442). G.A also thanks the Galileo Galilei Institute for the warm hospitality during part of the realization of this work.

References

- [1] P. J. Fox, R. Harnik, J. Kopp, and Y. Tsai, “Missing Energy Signatures of Dark Matter at the LHC,” *Phys.Rev.*, vol. D85, p. 056011, 2012, 1109.4398.
- [2] *et al.*, “Dark Matter Results from 225 Live Days of XENON100 Data,” 2012, 1207.5988.
- [3] R. Catena and P. Ullio, “A novel determination of the local dark matter density,” *JCAP*, vol. 1008, p. 004, 2010, 0907.0018.
- [4] R. Catena and P. Ullio, “The local dark matter phase-space density and impact on WIMP direct detection,” *JCAP*, vol. 1205, p. 005, 2012, 1111.3556.
- [5] F. Iocco, M. Pato, G. Bertone, and P. Jetzer, “Dark Matter distribution in the Milky Way: microlensing and dynamical constraints,” *JCAP*, vol. 1111, p. 029, 2011, 1107.5810.
- [6] S. Garbari, J. I. Read, and G. Lake, “Limits on the local dark matter density,” *Mon.Not.Roy.Astron.Soc.*, vol. 416, pp. 2318–2340, 2011, 1105.6339.
- [7] M. Pato, O. Agertz, G. Bertone, B. Moore, and R. Teyssier, “Systematic uncertainties in the determination of the local dark matter density,” *Phys.Rev.*, vol. D82, p. 023531, 2010, 1006.1322.
- [8] P. Salucci, F. Nesti, G. Gentile, and C. Martins, “The dark matter density at the Sun’s location,” *Astron.Astrophys.*, vol. 523, p. A83, 2010, 1003.3101.
- [9] W. de Boer and M. Weber, “The Dark Matter Density in the Solar Neighborhood reconsidered,” *JCAP*, vol. 1104, p. 002, 2011, 1011.6323.
- [10] S. H. Hansen, B. Moore, M. Zemp, and J. Stadel, “A Universal velocity distribution of relaxed collisionless structures,” *JCAP*, vol. 0601, p. 014, 2006, astro-ph/0505420.
- [11] C. Strege, R. Trotta, G. Bertone, A. H. Peter, and P. Scott, “Fundamental statistical limitations of future dark matter direct detection experiments,” *Phys.Rev.*, vol. D86, p. 023507, 2012, 1201.3631.
- [12] A. Djouadi *et al.*, “The Minimal supersymmetric standard model: Group summary report,” 1998, hep-ph/9901246.
- [13] G. Aad *et al.*, “Search for squarks and gluinos using final states with jets and missing transverse momentum with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions,” 2011, 1109.6572.
- [14] S. Chatrchyan *et al.*, “Search for Supersymmetry at the LHC in Events with Jets and Missing Transverse Energy,” *Phys.Rev.Lett.*, vol. 107, p. 221804, 2011, 1109.2352.
- [15] A. Djouadi, J.-L. Kneur, and G. Moultaka, “SuSpect: A Fortran code for the

- supersymmetric and Higgs particle spectrum in the MSSM,” *Comput.Phys.Commun.*, vol. 176, pp. 426–455, 2007, hep-ph/0211331.
- [16] G. Aad *et al.*, “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC,” *Phys.Lett.*, vol. B716, pp. 1–29, 2012, 1207.7214.
 - [17] P. P. Giardino, K. Kannike, M. Raidal, and A. Strumia, “Is the resonance at 125 GeV the Higgs boson?,” 2012, 1207.1347.
 - [18] D. Carmi, A. Falkowski, E. Kuflik, T. Volansky, and J. Zupan, “Higgs After the Discovery: A Status Report,” 2012, 1207.1718.
 - [19] H. E. Haber, “Challenges for nonminimal Higgs searches at future colliders,” 1995, hep-ph/9505240.
 - [20] H. Baer, V. Barger, and A. Mustafayev, “Implications of a 125 GeV Higgs scalar for LHC SUSY and neutralino dark matter searches,” *Phys.Rev.*, vol. D85, p. 075010, 2012, 1112.3017.
 - [21] A. Arbey, M. Battaglia, A. Djouadi, F. Mahmoudi, and J. Quevillon, “Implications of a 125 GeV Higgs for supersymmetric models,” *Phys.Lett.*, vol. B708, pp. 162–169, 2012, 1112.3028.
 - [22] L. J. Hall, D. Pinner, and J. T. Ruderman, “A Natural SUSY Higgs Near 126 GeV,” *JHEP*, vol. 1204, p. 131, 2012, 1112.2703.
 - [23] A. Arbey, M. Battaglia, A. Djouadi, and F. Mahmoudi, “The Higgs sector of the phenomenological MSSM in the light of the Higgs boson discovery,” 2012, 1207.1348.
 - [24] G. Aad *et al.*, “Search for neutral MSSM Higgs bosons decaying to $\tau^+\tau^-$ pairs in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector,” *Phys.Lett.*, vol. B705, pp. 174–192, 2011, 1107.5003.
 - [25] S. Chatrchyan *et al.*, “Search for neutral Higgs bosons decaying to tau pairs in pp collisions at $\sqrt{s}=7$ TeV,” *Phys.Lett.*, vol. B713, pp. 68–90, 2012, 1202.4083.
 - [26] P. P. Giardino, K. Kannike, M. Raidal, and A. Strumia, “Reconstructing Higgs boson properties from the LHC and Tevatron data,” *JHEP*, vol. 1206, p. 117, 2012, 1203.4254.
 - [27] T. Moroi, “The Muon anomalous magnetic dipole moment in the minimal supersymmetric standard model,” *Phys.Rev.*, vol. D53, pp. 6565–6575, 1996, hep-ph/9512396.
 - [28] L. Calibbi, R. Hodgkinson, J. Jones Perez, A. Masiero, and O. Vives, “Flavour and Collider Interplay for SUSY at LHC7,” *Eur.Phys.J.*, vol. C72, p. 1863, 2012, 1111.0176.
 - [29] E. Lunghi and J. Matias, “Huge right-handed current effects in $B \rightarrow K^*(K\pi)l^+l^-$ in supersymmetry,” *JHEP*, vol. 0704, p. 058, 2007, hep-ph/0612166.
 - [30] F. Mahmoudi, “SuperIso: A Program for calculating the isospin asymmetry of $B \rightarrow K^*\gamma$ in the MSSM,” *Comput.Phys.Commun.*, vol. 178, pp. 745–754, 2008, 0710.2067.
 - [31] F. Mahmoudi, “SuperIso v2.3: A Program for calculating flavor physics observables in Supersymmetry,” *Comput.Phys.Comm.*, vol. 180, pp. 1579–1613, 2009, 0808.3144.
 - [32] S. Chatrchyan *et al.*, “Search for $B_s^0 \rightarrow \mu^+\mu^-$ and $B^0 \rightarrow \mu^+\mu^-$ decays,” *JHEP*, vol. 1204, p. 033, 2012, 1203.3976.
 - [33] R. Aaij *et al.*, “Strong constraints on the rare decays $B_s \rightarrow \mu^+\mu^-$ and $B^0 \rightarrow \mu^+\mu^-$,” *Phys.Rev.Lett.*, vol. 108, p. 231801, 2012, 1203.4493.
 - [34] P. Gondolo *et al.*, “DarkSUSY: Computing supersymmetric dark matter properties

- numerically,” *JCAP*, vol. 0407, p. 008, 2004, astro-ph/0406204.
- [35] A. Hryczuk, R. Iengo, and P. Ullio, “Relic densities including Sommerfeld enhancements in the MSSM,” *JHEP*, vol. 1103, p. 069, 2011, 1010.2172.
 - [36] N. Arkani-Hamed, A. Delgado, and G. Giudice, “The Well-tempered neutralino,” *Nucl.Phys.*, vol. B741, pp. 108–130, 2006, hep-ph/0601041.
 - [37] J. R. Ellis, K. A. Olive, Y. Santoso, and V. C. Spanos, “Update on the direct detection of supersymmetric dark matter,” *Phys.Rev.*, vol. D71, p. 095007, 2005, hep-ph/0502001.
 - [38] M. Drees and M. M. Nojiri, “New contributions to coherent neutralino - nucleus scattering,” *Phys.Rev.*, vol. D47, pp. 4226–4232, 1993, hep-ph/9210272.
 - [39] M. Drees and M. Nojiri, “Neutralino - nucleon scattering revisited,” *Phys.Rev.*, vol. D48, pp. 3483–3501, 1993, hep-ph/9307208.
 - [40] J. Binney and S. Tremaine, “Galactic Dynamics: Second Edition,” *Princeton University Press*, 2008.
 - [41] H. Freudenreich, “Cobe’s galactic bar and disk,” *Astrophys.J.*, vol. 492, pp. 495–510, 1998, astro-ph/9707340.
 - [42] T. M. Dame, “The distribution of neutral gas in the milky way,” *AIP Conference Proceedings*, vol. 278, no. 1, pp. 267–278, 1992.
 - [43] J. F. Navarro, E. Hayashi, C. Power, A. Jenkins, C. S. Frenk, *et al.*, “The Inner structure of Lambda-CDM halos 3: Universality and asymptotic slopes,” *Mon.Not.Roy.Astron.Soc.*, vol. 349, p. 1039, 2004, astro-ph/0311231.
 - [44] A. W. Graham, D. Merritt, B. Moore, J. Diemand, and B. Terzic, “Empirical Models for Dark Matter Halos. III. The Kormendy relation and the $\log(\rho_{he}) - \log(R_e)$ relation,” *Astron.J.*, vol. 132, pp. 2711–2716, 2006, astro-ph/0608614.
 - [45] G. Bryan and M. Norman, “Statistical properties of x-ray clusters: Analytic and numerical comparisons,” *Astrophys.J.*, vol. 495, p. 80, 1998, astro-ph/9710107.
 - [46] E. Komatsu *et al.*, “Seven-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation,” *Astrophys.J.Suppl.*, vol. 192, p. 18, 2011, 1001.4538.
 - [47] X. Xue *et al.*, “The Milky Way’s Circular Velocity Curve to 60 kpc and an Estimate of the Dark Matter Halo Mass from Kinematics of 2400 SDSS Blue Horizontal Branch Stars,” *Astrophys.J.*, vol. 684, pp. 1143–1158, 2008, 0801.1232.
 - [48] Y. Akrami, C. Savage, P. Scott, J. Conrad, and J. Edsjo, “How well will ton-scale dark matter direct detection experiments constrain minimal supersymmetry?,” *JCAP*, vol. 1104, p. 012, 2011, 1011.4318.
 - [49] G. Bertone, D. G. Cerdeno, M. Fornasa, R. Ruiz de Austri, C. Strece, *et al.*, “Global fits of the cMSSM including the first LHC and XENON100 data,” *JCAP*, vol. 1201, p. 015, 2012, 1107.1715.
 - [50] D. G. Cerdeno, M. Fornasa, J.-H. Huh, and M. Peiro, “Nuclear uncertainties in the spin-dependent structure functions for direct dark matter detection,” 2012, 1208.6426.
 - [51] Y. Akrami, C. Savage, P. Scott, J. Conrad, and J. Edsjo, “Statistical coverage for supersymmetric parameter estimation: a case study with direct detection of dark matter,” *JCAP*, vol. 1107, p. 002, 2011, 1011.4297.

- [52] R. Trotta, F. Feroz, M. P. Hobson, L. Roszkowski, and R. Ruiz de Austri, “The Impact of priors and observables on parameter inferences in the Constrained MSSM,” *JHEP*, vol. 0812, p. 024, 2008, 0809.3792.
- [53] G. Bertone, D. G. Cerdeno, M. Fornasa, R. R. de Austri, and R. Trotta, “Identification of Dark Matter particles with LHC and direct detection data,” *Phys.Rev.*, vol. D82, p. 055008, 2010, 1005.4280.
- [54] G. B. Gelmini, P. Gondolo, A. Soldatenko, and C. E. Yaguna, “Direct detection of neutralino dark matter in non- standard cosmologies,” *Phys. Rev.*, vol. D76, p. 015010, 2007, hep-ph/0610379.
- [55] B. S. Acharya, G. Kane, S. Watson, and P. Kumar, “A Non-thermal WIMP Miracle,” *Phys. Rev.*, vol. D80, p. 083529, 2009, 0908.2430.
- [56] G. Arcadi and P. Ullio, “Accurate estimate of the relic density and the kinetic decoupling in non-thermal dark matter models,” *Phys.Rev.*, vol. D84, p. 043520, 2011, 1104.3591.
- [57] R. Catena, N. Fornengo, A. Masiero, M. Pietroni, and F. Rosati, “Dark matter relic abundance and scalar - tensor dark energy,” *Phys.Rev.*, vol. D70, p. 063519, 2004, astro-ph/0403614.
- [58] R. Catena, N. Fornengo, A. Masiero, M. Pietroni, and M. Schelke, “Enlarging mSUGRA parameter space by decreasing pre-BBN Hubble rate in Scalar-Tensor Cosmologies,” *JHEP*, vol. 0810, p. 003, 2008, 0712.3173.
- [59] G. Aad *et al.*, “Expected Performance of the ATLAS Experiment - Detector, Trigger and Physics,” 2009, 0901.0512.
- [60] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer, and T. Stelzer, “MadGraph 5 : Going Beyond,” *JHEP*, vol. 1106, p. 128, 2011, 1106.0522.
- [61] W. Beenakker, R. Hopker, and M. Spira, “PROSPINO: A Program for the production of supersymmetric particles in next-to-leading order QCD,” 1996, hep-ph/9611232.
- [62] G. Aad *et al.*, “Search for direct production of charginos and neutralinos in events with three leptons and missing transverse momentum in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector,” 2012, 1208.3144.
- [63] G. Aad *et al.*, “Search for direct slepton and gaugino production in final states with two leptons and missing transverse momentum with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV,” 2012, 1208.2884.
- [64] G. Aad *et al.*, “Search for top and bottom squarks from gluino pair production in final states with missing transverse energy and at least three b-jets with the ATLAS detector,” 2012, 1207.4686.
- [65] G. Aad *et al.*, “Search for direct top squark pair production in final states with one isolated lepton, jets, and missing transverse momentum in $\sqrt{s} = 7$ TeV pp collisions using 4.7 fb⁻¹ of ATLAS data,” 2012, 1208.2590.
- [66] G. Aad *et al.*, “Search for a supersymmetric partner to the top quark in final states with jets and missing transverse momentum at $\sqrt{s} = 7$ TeV with the ATLAS detector,” 2012, 1208.1447.
- [67] “Search for dark matter candidates and large extra dimensions in events with a jet and missing transverse momentum with the atlas detector,” Tech. Rep. ATLAS-CONF-2012-084,

CERN, Geneva, Jul 2012.

- [68] “Search for scalar bottom pair production in Pnal states with missing transverse momentum and two b-jets in pp collisions at $\sqrt{s}=7$ tev with the atlas detector,” Tech. Rep. ATLAS-CONF-2012-106, CERN, Geneva, Aug 2012.
- [69] G. Aad *et al.*, “Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using 4.7 fb^{-1} of $\sqrt{s} = 7$ TeV proton-proton collision data,” 2012, 1208.0949.
- [70] “Search for squarks and gluinos with the atlas detector using final states with jets and missing transverse momentum and 5.8 fb^{-1} of $\sqrt{s}=8$ tev proton-proton collision data,” Tech. Rep. ATLAS-CONF-2012-109, CERN, Geneva, Aug 2012.
- [71] H. Baer, V. Barger, A. Lessa, W. Sreethawong, and X. Tata, “Wh plus missing- E_T signature from gaugino pair production at the LHC,” *Phys.Rev.*, vol. D85, p. 055022, 2012, 1201.2949.
- [72] J. A. Conley, J. S. Gainer, J. L. Hewett, M. P. Le, and T. G. Rizzo, “Supersymmetry Without Prejudice at the 7 TeV LHC,” *Physical Review D*, 2011, 1103.1697.
- [73] H. K. Dreiner, M. Kramer, and J. Tattersall, “How low can SUSY go? Matching, monojets and compressed spectra,” *Europhys.Lett.*, vol. 99, p. 61001, 2012, 1207.1613.
- [74] G. Belanger, M. Heikinheimo, and V. Sanz, “Model-Independent Bounds on Squarks from Monophoton Searches,” *JHEP*, vol. 1208, p. 151, 2012, 1205.1463.
- [75] G. Aad *et al.*, “Search for scalar bottom pair production with the ATLAS detector in pp Collisions at $\sqrt{s} = 7$ TeV,” *Phys.Rev.Lett.*, vol. 108, p. 181802, 2012, 1112.3832.
- [76] G. Aad *et al.*, “Search for light top squark pair production in final states with leptons and b-jets with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions,” 2012, 1209.2102.
- [77] A. Choudhury and A. Datta, “Many faces of low mass neutralino dark matter in the unconstrained MSSM, LHC data and new signals,” *JHEP*, vol. 1206, p. 006, 2012, 1203.4106.
- [78] O. Adriani *et al.*, “PAMELA results on the cosmic-ray antiproton flux from 60 MeV to 180 GeV in kinetic energy,” *Phys.Rev.Lett.*, vol. 105, p. 121101, 2010, 1007.0821.
- [79] C. Weniger, “A Tentative Gamma-Ray Line from Dark Matter Annihilation at the Fermi Large Area Telescope,” *JCAP*, vol. 1208, p. 007, 2012, 1204.2797.
- [80] W. Buchmuller and M. Garny, “Decaying vs Annihilating Dark Matter in Light of a Tentative Gamma-Ray Line,” *JCAP*, vol. 1208, p. 035, 2012, 1206.7056.
- [81] G. Aad *et al.*, “Search for dark matter candidates and large extra dimensions in events with a jet and missing transverse momentum with the ATLAS detector,” 2012, 1210.4491.
- [82] A. Choudhury and A. Datta, “New limits on top squark NLSP from LHC 4.7 fb^{-1} data,” 2012, 1207.1846.